



International Journal of Fisheries and Aquatic Studies

ISSN: 2347-5129
IJFAS 2014; 2(2): 130-136
© 2013 IJFAS
www.fisheriesjournal.com
Received: 11-09-2014
Accepted: 30-09-2014

Benjamin J. Gonzales
*College of Fisheries and Aquatic
Sciences, Western Philippines
University, Puerto Princesa
Campus, Santa Monica,
Palawan, 5300*

Roger G. Dolorosa
*College of Fisheries and Aquatic
Sciences, Western Philippines
University Puerto Princesa
Campus, Santa Monica,
Palawan, 5300.*

Honorio B. Pagliawan
*College of Fisheries and Aquatic
Sciences, Western Philippines
University Puerto Princesa
Campus, Santa Monica,
Palawan, 5300.*

Maria Mojena G. Gonzales
*College of Fisheries and Aquatic
Sciences, Western Philippines
University Puerto Princesa
Campus, Santa Monica,
Palawan, 5300.*

Correspondence:
Maria Mojena G. Gonzales
*College of Fisheries and Aquatic
Sciences, Western Philippines
University Puerto Princesa
Campus, Santa Monica,
Palawan, 5300*

Marine resource assessment for sustainable management of Apulit Island, West Sulu Sea, Palawan, Philippines

**Benjamin J. Gonzales, Roger G. Dolorosa, Honorio B. Pagliawan and Maria
Mojena G. Gonzales**

Abstract

The marine resources in Apulit Island, Palawan were assessed to evaluate its status after decades of protection. The hard coral cover in six sites ranged from 6.88 to 71.25%. Some reefs have been overgrown with macroalgae. The reef fish biomass and density were relatively high, but poor in species diversity. Subsequent underwater footage and anecdotal information revealed a large change of fish assemblage, which may have occurred during the transition period from previous owner/operator to the recent owner/operator: This situation may highlight the role of local government in the management of coastal waters and its resources.

Keywords: Marine Protected Area (MPA), Coral cover, Fish assemblage

1. Introduction

Apulit is an Island resort located at the center of Taytay Bay, Palawan. As excellent dive site, the Resort was established in early 90's by private entity. This was accompanied by municipal ordinance declaring the island as marine protected area (MPA). Well managed MPA can help sustain and enhance the populations of target species within and outside the protected areas [1, 2].

Success of several community-based MPAs in the Philippines were reported by Alcala [3] and White *et al.* [4]. However, there is a concern that the majority of the MPAs are not effectively managed [5, 4] and are mostly not achieving marine biodiversity conservation [6], fisheries enhancement or other objectives [7].

Improved habitat conditions in terms of coral cover and fishery yields associated with MPAs are associated with management success of the area [8]. Thus this study aims to assess the marine resources of Apulit Island in terms of coral cover and reef associated fish and invertebrates which can be used as management success indicator.

2. Materials and Methods

2.1 The study site

Apulit Island is situated in the eastern coast of Northern Palawan more specifically located 10°57.237' North and 119°36.824' East (Fig. 1). It is part of Barangay Mayteged, Municipality of Taytay, Province of Palawan. It is 17 kilometers north-east of the municipality of Taytay. Apulit Island is predominantly sloping and mountainous with its highest peak towering at 164 meters above sea level.

2.2 Survey method for coral cover

Transect method modified after Hodgson *et al.* [9] was used to evaluate the present state of the coral reefs around Apulit Island and its nearby vicinity. This method counts at family level of some fish and invertebrate species typically targeted as food species or collected for curio trade and substrate categories chosen to detect large scale changes were made.

Selected six sites for permanent monitoring were based on the suggestions of the Resort's staff knowledgeable of the area. Five of these were within the jurisdiction of Apulit Island and the other one is in a nearby island. A 100 m transect was established per site except in Goose neck (Transect 5) where only a 50 m transect was laid due to its depth. Coordinates were recorded using Garmin II GPS (Fig. 2).




Fig 1: Map of Palawan showing the location of the study site, Apulit Island in Taytay Bay, Palawan Philippines.




Fig 2: Transects along Apulit Island for reef check

2.3 Fish visual census

The same stations for reef check were used for this fish visual census (Table 1 and Fig. 2) following the method of English *et al.* [10]. The survey used three 30m-transect lines per station.

Fish species encountered along the transect lines were identified up to the species level. The total lengths of fish were estimated in centimeters, which were used to estimate biomass. Parameters *a* and *b* were taken from Kulbicki *et al.* [11]. References used for fish identification were: Gonzales [12], Fishbase [13], Kuitert and Debelius [14], Randall [15], and Masuda *et al.* [16].

3. Results and Discussion

3.1 Coral reef

Percent hard coral (HC) cover ranged from 6.88 - 71.25% (Table 1). The site (T4, Lopez Reef) with the highest (71.25%) live coral cover distantly located from Apulit Island while an area close to the resort (T1; Platform) had the lowest 6.88% coral cover. The data also suggest an inverse relationship between live coral cover and macroalgae (Fig 3). Turbid waters were drained towards T1 and siltation must have reached up to the Noah's Rock (T3) where HC cover is only 33%. Analyzing the report of WWF [17], would reveal about 5% hard coral cover in this site. Turbid water can have stressful effects on corals while nutrients that are drained in the area could enhance the growth of algae which outgrow and compete with corals for space and sunlight. In Apulit Island, percent composition of HC is inversely proportional ($r^2 = -0.7163$) with macroalgae (Fig. 3).

Table 1: Condition of reefs in Apulit Island, Honda Bay, Puerto Princesa City, Palawan, Philippines, October, 2006.

Station	Total coral cover (%)	Reef Condition*
1	6.88	Poor
2	54.38	Good
3	33.12	Fair
4	71.25	Good
5	41.25	Fair
6	33.12	Fair
Average	40	Fair

*Adjectival rating on coral reef condition was used to assess the health of coral community: <25% - poor; >25% but <50% - fair; >50% but <75 - good and >75 - excellent. Gomez *et al.* [20]

There are no available baseline data prior to the establishment of MPA and resort in Apulit Island. Hence, this study acts as a baseline data for management of its resources. It has been noted also that areas close to Apulit Island (T1, T3, T6) has lower percent HC cover compared to those that are a bit far (T4, T2, T5). For T6 to have low coral cover is understandable because although it is close to the island yet prone to fishing as it is found at the eastern side and is not visible from the resort. Evidence of continued fishing in T5 is reflected in the state of some fragmented stag horn coral colonies and the scarcity of commercially

important fish and invertebrate species. Unsustainable or damaging fishing practices could damage the coral reef and as an effect could degrade fish catch of fishermen [18].

In early 90's, there were reports on effect of climate change and illegal fishing activity in coral reef within Palawan specifically in Tubbataha Reefs [19]. Recovery of degraded reefs is slow or will never occur naturally once overgrown with algae. In some sites, macroalgae have flourished and have taken over the reef area. For example, the base of branching *Acropora* sp. close to the resort had already been colonized by *Sargassum* spp. In many stations, it has been noted that small colonies of corals are already being covered with *Sargassum* spp and other species of macroalgae. We foresee that these reefs will become a *Sargassum* bed.




Fig 3: Relationship between percent hard coral cover and nutrient indicator algae.

Invertebrate survey along the established permanent transects indicated that areas around Apulit Island and its vicinity has been overfished. For example, although T4 had the highest percent HC cover yet there is not much reef associated invertebrates. Only the unexploited *Diadema* sp. or tayom for the locals are quite abundant in most areas.

In T4, boring Tridacnid clams *Tridacna crocea* are abundant with a density of about 31 ind./100 m². This is because the area is shallow and has high percent live hard coral cover of massive and sub massive forms. Burrowing *Tridacna crocea* seems to be not exploited may be because of their small size and the difficulty in extracting the meat. Several non-boring giant clams like *Hippopus porcellanus* and *Tridacna squamosa* species were found in T1 which is again inaccessible to fishing.

The average density (6.81 ind./100 m²) (Table 2) of giant clams in sampling areas was a bit higher compared to that in Caramay (0.7 ind./100 m²) and Green Island (1.5 ind./100 m²) as reported by Condesa [21] but comparable to those from Tubbataha Reefs (4.4 ind./100 m²) [22] and Malampaya Sound, Taytay, Palawan (5.99 ind./m²) [23]. However, the absence of high valued sea cucumbers and the low densities of commercially important coral associated invertebrates is indicating a recovering population.

Table 2: Invertebrates density (ind./100 m²) recorded at different transects around and near Apulit Island, Taytay, Palawan.

Invertebrates	T1	T2	T3	T4	T5	T6	Average
Balatan (Holothurians)					0.4		0.07
Taklobo (Tridacnids)	4.2	2.8	0.25	30.67	1.2	1.75	6.81
Tayom (Diadema sp.)	37.4	38.4	14.75	126.67	10.4	2.25	38.31
Tirik (Sea urchin)	1.2	2.8	0.5		1.6	2	1.35

3.2 Fish census

The reef fishes around Apulit Island can be classified as having high biomass (wt), highly diverse (no. of species), and high in density (no. of individual/unit area) except the Lopez Island reef which is considered as unprotected area.

The total number of species observed in six stations was 110 (Appendix A), out of this 99 were identified to species level belonging to 20 families (Table 3). The families with high number of species were Pomacentridae (Damsel), (30), Labridae (Wrasses), (25), Scaridae (Parrotfish), (12), and Serranidae (Grouper) and Chaetodontidae (Butterflyfish), (6). The total number of commercially valuable species was 53, comprising 48.38% of the total number of species observed along the transect lines. In this report, commercially

important species include fishes belonging to families of Labridae (Wrasses) (47.17%); Scaridae (Parrot fishes), (22.64%); Serranidae (Groupers), (11.32%); Nemipteridae (Breems), (7.55%) Lutjanidae (Snappers), (5.66%); Caesionidae (fusiliers) (3.77%); and Acanthuridae (Surgeonfishes), (1.87%).

The number of chaetodons (Butterfly fishes) was considerably low (5.35%). Butterfly fishes have been used as indicators for reef health since they are highly associated with coral reef. Pomacentridae (Damsel fishes) is known to be highly territorial reef fish. In this survey, it was noted that this group appears moderately high in abundance (26.75%). The estimated species diversity was 0.108 species/m².

Table 3: Family composition and species distribution in six stations.

	Family	Goose Breast	Goose Neck	Stations Noah's Rock	Lopez Is.	Eastside	Platform	Total
1	Acanthuridae	1	1				2	4
2	Apogonidae	2	2		1			5
3	Caesionidae	1	2				1	4
4	Carangidae	1	1				1	3
5	Chaetodontidae	3	2	2	3	2	1	13
6	Ephippidae	1					1	4
7	Holocentridae	1				1		2
8	Labridae	6	13	10	8	12	9	58
9	Lethrinidae	1	1	2			1	5
10	Lutjanidae	1	2	2	21	1	3	11
11	Mullidae					1		1
12	Nemipteridae	1		2	3			6
13	Pinguipedidae					1		2
14	Pomacanthidae	1		1	1	1	1	5
15	Pomacentridae	9	8	22	9	12	5	65
16	Pseudochromidae	1	1	1	1	1	1	7
17	Scaridae	2	1	4	4	3	1	14
18	Serranidae	5	3	3	2	2	8	23
19	Siganidae	3	1	3	1	1	2	11
20	Tetraodontidae	1		1				2

The total fish biomass was estimated at 371.65 mt/km². Noah's Rock has the highest biomass, 132.29 mt/km² followed by Platform area (99.13 mt/km²) (Table 4). The high biomass in these two sites may be attributed to presence

of large sizes of target species (e.g. Groupers, Snappers, Sweetlips, and Caranx). This high biomass fish can be also attributed to the proper management and effective protection of the island.

Table 4: Profile of associated reef fish species and some ecological indicators of reef fish conditions in six sampling stations around Apulit Island.

Station	Depth (m)	Families (species/ 1,000 m ²)	Diversity (individual/ 1000 m ²)	Density (mt/km ²)	*Biomass
Noah's Rock	5 – 6	13	50	1,171	132.29
Platform	5 – 6	13	37	2,035	99.13
Gooseneck	5 – 13	13	38	1,864	66.67
Goose breast	3 – 5	18	41	1,173	40.80
Eastside	2 – 5	13	41	1,420	16.51
Lopez Island	2 – 3	11	34	617	16.27

*mt = metric tons **Total Fish Biomass** = 371.65 mt/km² **Average Biomass** = 61.94 mt/km²

Of the six stations surveyed, the Noah's Rock station had the most diverse fish species of 50 species/1,000 m² followed by Goose breast and Eastside Stations having 41 species/1,000 m². Platform area has 37 species/1,000 m² (17.67%). On the hand, Lopez Island rock was the least diverse among the six sampling stations, 34 species/1,000 m².

Based on the category of Hilomen *et al.*^[24], the average fish biomass (61.94 mt/km²) of Apulit Island was categorized as high, compared to St. Paul Bay (29.26 mt/km²)^[12], Honda Bay (27.11 mt/km²)^[25], and Tabuyo Fish Sanctuary in

Caramay, Roxas, (22.94 mt/km²)^[26], which were all moderate in terms of fish biomass. This result was indicative that reefs around Apulit Island were healthy. A healthy Philippine reef produces 5-37 mt/km² of fish per year^[27].

The average fish density in the island estimated at 1.38 indv/m² or 1,380.00 indvs/1000 m² was categorized as moderate in terms of abundance. The high category ranged from 2268 to 7592 indv/1000 m² density^[24].

Five stations have poor diversity 34-41 species/ 1000 m², while only one station (the Noah's Rock) was categorized as moderately diverse, having 50 species/ 1000 m².

Appendix

Fish species observed in Apulit Island Taytay, Palawan

SI. No.	Species Name	SI. No.	Species Name	SI. No.	Species Name
1	<i>Acanthochromis polyacanthus</i>	38	<i>Choerodon anchorago</i>	74	<i>Platax tiera</i>
2	<i>Acanthurus pyroferus</i>	39	<i>Chromis viridis</i>	75	<i>Plectroglyphidodon jonhstonianus</i>
3	<i>Acanthurus striatus</i>	40	<i>Chrysiptera cyanea</i>	76	<i>Plectroglyphidodon lacrymatus</i>
4	<i>Acanthurus xanthopterus</i>	41	<i>Chrysiptera tricincta</i>	77	<i>Plectropomus leopardus</i>
5	<i>Amblyglyphidodon aureus</i>	42	<i>Cirrhitilabrus cyanopleura</i>		<i>Plectropomus maculatus</i>
6	<i>Amblyglyphidodon curacao</i>	43	<i>Coris aurineata</i>	78	<i>Plectropomus oligacanthus</i>
7	<i>Amblyglyphidodon leucogaster</i>	44	<i>Dascyllus reticulatus</i>	79	<i>Pomacentrus sp. (palata)</i>
8	<i>Amphiprion clarkii</i>	45	<i>Dascyllus trimaculatus</i>	80	<i>Pomacentrus alexanderea</i>
9	<i>Anampses geographicus</i>	46	<i>Diproctacanthus xanthurus</i>	81	<i>Pomacentrus amboinensis</i>
10	<i>Anampses caeruleopunctatus</i>	47	<i>Dischistodus melanotus</i>	82	<i>Pomacentrus lividus</i>
11	<i>Anyperodon leucogrammicus</i>	48	<i>Dischistodus perspicillatus</i>	83	<i>Pomacentrus molucensis</i>
12	<i>Apogon angustatus</i>	49	<i>Epibulus insidiator</i>	84	<i>Pomacentrus nagasakiensis</i>
13	<i>Apogon sp. (yg)</i>	50	<i>Epinephelus fasciatus</i>	85	<i>Pomacentrus rhodontus</i>
14	<i>Apogon trimaculatus</i>	51	<i>Epinephelus fuscoguttatus</i>	86	<i>Pomacentrus richardsoni</i>
15	<i>Arothron nigropunctatus</i>	52	<i>Epinephelus qouyanus</i>	87	<i>Pomacentrus sp.</i>
16	<i>Atule mate</i>	53	<i>Glyphidodontops parasema</i>	88	<i>Pomacentrus sp. (binduyan)</i>
17	<i>Bodianus mesothorax</i>	54	<i>Halichoeres chloropterus</i>	89	<i>Pomacentrus sp. 2</i>
18	<i>Bodianus perditio</i>	55	<i>Halichoeres hoeveni</i>	90	<i>Pomacentrus vauli</i>
19	<i>Bolbometopon bicolor</i>	56	<i>Hemiglyphidodon plagiometopon</i>	91	<i>Pomachromis richarsoni</i>
20	<i>Caesio cunning</i>	57	<i>Hemigymnus fasciatus</i>	92	<i>Psuedochromis fuscus</i>
21	<i>Canthigaster coronata</i>	58	<i>Hemigymnus melaterus</i>	93	<i>Pterocaesio tile</i>
22	<i>Carangoides plagiotaenae</i>	59	<i>Henicopus varius</i>	94	<i>Scarus sp. (brown)</i>
23	<i>Caranx ignobilis</i>	60	<i>Labracinus cyclophthalmus</i>	95	<i>Scarus sp. (green)</i>
24	<i>Centropyge multifasciatus</i>	61	<i>Labroides demidiatus</i>	96	<i>Scarus sp. (blue)</i>
25	<i>Centropyge vroliki</i>	62	<i>Labropsis manabei</i>	97	<i>Scarus sp. (gray)</i>
26	<i>Cephalopholis boenak</i>	63	<i>Lethrinus sp.</i>	98	<i>Scolopsis bilineatus</i>
27	<i>Cephalopholis cyanostigma</i>	64	<i>Lutjanus carponotatus</i>	99	<i>Scolopsis ciliatus</i>
28	<i>Cetoscarus bicolor (juv)</i>	65	<i>Lutjanus decussatus</i>	100	<i>Scolopsis margaritifera</i>
29	<i>Chaetodon baronesa</i>	66	<i>Lutjanus ornatus</i>	101	<i>Scolopsis dubiosus</i>
30	<i>Chaetodon lineolatus</i>	66	<i>Lutjanus vittatus</i>	102	<i>Siganus corallinus</i>
31	<i>Chaetodon speculum</i>	67	<i>Myripristis sp.</i>	103	<i>Siganus punctatissimus</i>
32	<i>Cheilinus celebicus</i>	68	<i>Paraglyphidodon melas</i>	104	<i>Siganus virgatus</i>
33	<i>Cheilinus chlorurus</i>	69	<i>Paraglyphidodon negroris</i>	105	<i>Siganus vulpinus</i>
34	<i>Cheilinus diagrammus</i>	70	<i>Parapersis hexophthalma</i>	106	<i>Stethojulis bandanensis</i>
35	<i>Cheilinus fasciatus</i>	71	<i>Parupeneus indicus</i>	107	<i>Stethojulis trilineata</i>
36	<i>Chelmon rostratus</i>	72	<i>Pentapodus canius</i>	108	<i>Thalassoma lunari</i>
37	<i>Chlorurus sordidus</i>	73	<i>Platax orbicularis</i>		

4. Conclusion and Recommendation

Since conditions of Apulit Island coral reef cover and fish assemblage were better compared to other coral reefs in Palawan and the Philippines, its integrity should be maintained and reefs must be protected to further improve its status. For the purpose of proper eco-tourism planning and management, the Island and its reefs should be classified into

zones. Area or zones for biodiversity conservation and multi-use, including eco-tourism must be distinct and clearly understood. In this way policies and regulations would be clear not only at the Island management level, but also with the municipality.

Coral reef areas in the island should be managed through the establishment of marine protected area, where a fish

sanctuary should be established as a core (no take) zone to serve as reservoir and generator of marine plants and animals in the area, and multipurpose zone to cater to the needs of fishers and other stakeholders for their respective interest. The reefs should be managed in a way that it would not only generate and enhance the reef ecosystem, but also extend aesthetic and educational values to visitors.

Damaged coral reefs if properly protected could recover within the span of five to ten years (Veron, 1986). In addition, regular monitoring of the reefs shall be done in order to detect changes in reefs' conditions, so that managers could readily adjust management schemes parallel to the status and trends of the reefs, especially in relation to climate change.

In addition, it is recommended that growth of *Sargassum* (seaweeds) and other algae that cover coral colonies must be monitored and controlled to regulate its negative effects to the corals. Situating permanent mooring buoys should be provided in every dive/activity site to minimize coral and other marine life damage every time anchor hits the bottom. There should discourage feeding of fishes. Feeding fishes in the wild alters the natural capability of fish to survive in the wild. It weakens the ability of fish to find food by themselves.

5. Acknowledgement

We wish to acknowledge Mr. Keizaburo O. Homma and his officers and staff, the management of Club Noah Isabelle for allowing us the conduct of resource assessment in the island. Special thanks are due to our partners, Mr. Ricky Tamparong, the Manager of the resort, and all his staff for their hospitality, serving as guides and associates during the survey. We are obliged to Maria Victoria Matillano of WWF –Malampaya, Palawan in her valuable contribution and assistance in coordination, documentation, and night lectures.

6. References

- Sale PF, Cowen RK, Danilowicz BS, Jones GP, Kritzer JP, Lindeman KC *et al.* Critical science gaps impede use of no-take fishery reserves. *Trends Ecol Evol* 2005; 20:74–80.
- Russ GR. Yet another review of marine reserves as reef fishery management tools. In: Sale PF, editor. *Coral reef fishes: dynamics and diversity in a complex ecosystem*. Amsterdam: Academic Press, 2002.
- Alcala AC. Community-based coastal resource management in the Philippines: A case study. *Ocean and Coastal Management* 1998; 38:179-186.
- White AT, Aliño PA, Menesses A. Creating and managing marine protected areas in the Philippines. *Fisheries Improved for Sustainable Harvest Project*, Coastal Conservation and Education Foundation, Inc. and the University of the Philippines Marine Science Institute, Cebu City, Philippines, 2006, 83.
- Christie P, White A, Deguit E. Starting point or solution? Community-based marine protected areas in the Philippines. *Journal of Environmental management* 2002; 66:441-454.
- Weeks R, Russ GR, Alcala AC, White AT. Effectiveness of Marine Protected Areas in the Philippines for Biodiversity Conservation. *Conservation Biology*, Vol 10, No. 1523-1739. 2009 Society for Conservation Biology, 2009, 1-10.
- Maliao RJ, White AT, Maypa AP, Turigan RG. Trajectories and magnitude of change in coral reef fish populations in Philippine marine reserves: A meta-analysis. *Coral Reefs* 2009; 28:809-822.
- Maypa AP, White AT, Cañares E, Martinez R, Osorio RL, Aliño P *et al.* Marine Protected Area Management Effectiveness: Progress and Lessons in the Philippines in *Coastal Management* 2012; 40(5):510-524.
- Hodgson G, Kiene W, Mihaly J, Liebeler J, Shuman C, Maun L *et al.* Reef Check Instruction Manual: A Guide to Reef Check, Coral Reef Monitoring, Institute of the Environment, University of California at Los Angeles, 2004.
- English S, Wilkinson C, Baker V. Survey manual for tropical marine resources. Edn 2, Australian Institute of Marine Science, Townsville, Australia, 1997, 390.
- Kulbicki M, Guillemot N, Amand MA. General approach to length-weight relationships for New Caledonian Lagoon fishes. *Cybius* 2005; 29:235-252.
- Gonzales BJ. Food Fishes of Palawan. Edn 2, Bureau of Fisheries and Aquatic Resources-fisheries Resource Management Project-Philippine Information Agency, Quezon City, Philippines, 2000.
- Fish Base. World fish Center, Malaysia, 2004.
- Kuiter RH, Debelius H. Southeast Asia tropical fish guide. IKAN Unterwasserarchiv, 1997.
- Randall JE. Red Sea Reef Fishes. Immel Publishing Limited London, 1992.
- Matsuda H, Amaoka K, Araga C, Uyeno T, Yoshino T. The Fishes of the Japanese Archipelago, Tokai University Press, 1984.
- WWF International. No date. So what is a representative network of MPAs? (Available at www.panda.org/what_we_do/how_we_work/conservation/marine/protected_areas/increasing_protection/mpa_networks/).
- Dubinsky Z. Coral Reefs. Ecosystem of the World. Elsevier. New York, 1990, 1-550
- White AT, Calumpang H. Saving Tubbataha Reefs, Earthwatch Expedition to the Philippines. *Siliman Journal* 1993; 36(2):77-105
- Gomez ED, Alcala AC, San Diego AC. Status of Philippine coral reefs. Proceeding of the 4th International Coral Reef Symposium, Manila. Marine Science Center, University of the Philippines, Diliman, Quezon City, Philippines, 1981; 1:275-282.
- Condesa LD. Gleaning and abundance of commercially valuable intertidal invertebrates in Green Island and Caramay, Roxas, Palawan. Undergraduate thesis, Western Philippines University-Puerto Princesa Campus, Puerto Princesa City, Palawan, Philippines. 2005; 42.
- Sayson SM. Assessment of giant clams and recommendations for management in Binaluan Bay, Malampaya Sound, Taytay, Palawan, Philippines. Undergraduate thesis, Western Philippines University-Puerto Princesa Campus, Puerto Princesa City, Palawan, Philippines, 2003, 56.
- Dolorosa RG, Schoppe S. Focal benthic mollusks (Mollusca: Bivalvia and Gastropoda) of selected sites in Tubbataha Reef national Marine Park, Palawan, Philippines. *Science Diliman* 2005; 17(2):1-8.
- Hilomen VV. Associated reef fishes In *Manual on the Standardization of field methods for assessment of coral resources*. Sulu-Celebes Project (PCICDSCS), 1998.

25. Gonzales BJ, Galon W, Bicera J, Pagliawan HB, Rodriguez EF, Bactol GP *et al.* Resource Assessment for Sanctuary Establishment, Saint Paul Bay, Western Palawan, Philippines, Saguda Palawan, Inc. Puerto Princesa City, Palawan, 2005.
26. Pagliawan HB, Palla HP. Reef Fish communities of Tabuyo Fish Sanctuary, Caramay, Roxas, Palawan. Unpublished Report, 2005.
27. Alcala AC. Marine reserves in the Philippines: historical development, effects and influence on marine conservation policy. Bookmark, Makati City, 2001, 115.