



International Journal of Fisheries and Aquatic Studies

ISSN: 2347-5129

(ICV-Poland) Impact Value: 5.62

(GIF) Impact Factor: 0.352

IJFAS 2016; 4(1): 85-89

© 2016 IJFAS

www.fisheriesjournal.com

Received: 17-10-2015

Accepted: 21-11-2015

Muneesh kumar

Department of Zoology Govt.
S.S.L. Jain P.G. College, Vidisha,
Barkatullah University, Bhopal
(M.P.) India.

Mansa Ram

Department of Zoology Govt.
Degree College Bhaderwah,
University of Jammu, Jammu.
India.

Toxicity of some heavy metals on blood characteristics of freshwater fish *Clarias batrachus*

Muneesh kumar, Mansa Ram

Abstract

The environment is impacted by ongoing pollution, caused by both natural as well as environmental factors such as industrialization and mining. Heavy metals causes as major problem because they are toxic and tend to accumulate in living organisms. So the present study was conducted to determine the effects of sublethal concentrations of $ZnSO_4$, $CdCl_2$ and $CuSO_4$ as a pollutant on some haematological parameters of *Clarias batrachus* were exposed to three sublethal levels. Sublethal concentrations used were 6.5 ppm for cadmium, 4.6 ppm for Zn and 0.7 ppm for Cu. The results obtained revealed that there was significant difference in hematocrit, Exposure of *Clarias batrachus* to sublethal concentrations of Zn and Cd resulted in significant decrease ($p < 0.05$) in erythrocyte count and haemoglobin content, while there was a slight increase in case of copper. The microscopical observations showed that exposure of *Clarias batrachus* to sublethal concentrations of heavy metals 6.5 ppm for cadmium, 4.6 ppm for Zn and 0.7 ppm for Cu resulted in some changes in the morphology of the R.B.Cs which tended to lose its ellipsoidal shape acquiring different shapes (poikilocytosis or anisocytosis).

Keywords: Heavy metals, Hematological parameters, *Clarias batrachus*.

1. Introduction

Blood alterations in fish or damage of hemopoietic tissue organs may be associated either with changes in environmental conditions or with water borne pollutants (McCay, 1929^[23]; Dawson, 1935^[12] and Gradner and Yevich 1969b^[15]). Most of the earlier studies on the effect of intoxication with pollutants on blood characteristics were based on differential counts of W.B.C. and R.B.C., as well as the relative concentration of haemoglobin content and hematocrit values (Khadre, 1990; Amin, 1992^[7]; Houston *et al.*, 1993^[20]; Adak, 1995^[5] and Abdel Magid, 1997). In the present study morphological characteristics of red blood cells were studied beside the other haematological parameters on *C. batrachus* exposed to sublethal concentrations of some heavy metals (Cu, Zn and Cd).

2. Material and methods

Adult and live fish *C. batrachus* were collected from the fish farm Patra and Bhadbhada Bhopal M.P.) and were brought to the laboratory, cleaned by using 0.1% $KMnO_4$ to avoid dermal infection. Only healthy fishes (Length: 12-15cm, Weight: 50-60g) were taken for experiment. Fishes were acclimatized in glass aquaria for 15 days and were fed with fish food (earthworms) and water in the aquaria was replaced by freshwater at every 24h. They were put in glass aquaria (60 liters capacity) for about two weeks for acclimatization. Then the tested media were prepared. Experimental fish after this acclimatization were transferred one hour after preparation of the media, into the experimental aquaria.

Experimental glass aquaria were cleaned carefully and washed with water and acetone then with running tap water several times and finally.

Pollutants: 1000 ppm stock solutions of the three metals were prepared. Cadmium as cadmium chloride ($CdCl_2 \cdot H_2O$), zinc as zinc sulphate ($ZnSO_4 \cdot 7H_2O$) and copper as copper sulphate ($Cu SO_4 \cdot 5H_2O$). Standard solutions were stored in clean bottles and diluted to desired concentration just before the beginning of the experiment. Sublethal concentrations used were Zn (4.6 ppm), Cd (6.5 ppm) and Cu (0.7 ppm) (Abdel Barr 1997).

R.B.Cs count: R.B.Cs counts were done according to the method of Hesser (1960)^[17] and Wintrob (1967) using improved Neubauer haemocytometer. Haemoglobin content was determined according to the calorimetric method of Sahli (Hesser 1960^[17]).

Correspondence

Muneesh kumar

Department of Zoology Govt.
S.S.L. Jain P.G. College, Vidisha,
Barkatullah University, Bhopal
(M.P.) India.

3. Results

Normal red blood cells

The erythrocytes of normal fish are ellipsoidal in shape (the short axis nearly two thirds of the long axis). They are nucleated with a centrally located nucleus, which is ellipsoidal as well. The measurement of the cells varies between $11 \times 5 \mu\text{m}$ and $7 \times 6 \mu\text{m}$ with a mean of $8.8 \pm 1.4 \times 5.7 \pm 0.9 \mu\text{m}$ for long and short axis. The nuclei are dark purple in colour and measure between $4 \times 3.5 \mu\text{m}$ and $5 \times 3 \mu\text{m}$ with a mean of $4.2 \pm 0.4 \mu\text{m} \times 3.3 \pm 0.5 \mu\text{m}$ on the long and short axis respectively. R.B.Cs count in normal fish ranged from 420000 to 720000 cell/ mm^3 with a mean value 445000 ± 250000 cell/ mm^3 .

Haemoglobin content

Haemoglobin content was found to vary between 1.38 and 1.54/ 100 ml blood for normal fish of *C. batrachus*.

(I) Morphological variations in red blood cells

(1) Effect of sublethal concentration of Zinc (4.6 ppm)

Lysis of red blood cells is quite clear upon examination of the blood films after 4 days of exposure to 4.6 ppm Zn (cell wall of erythrocyte disintegrated and the nuclear material were clumped) (Fig. 1a). On the other hand diameter of the red blood cells showed a non-significant decrease ($8.1 \pm 1.2 \mu\text{m}$ in treated fish and $8.8 \pm 1.4 \mu\text{m}$ in control, $t = -2.0$ & $P < 0.5$). After 8 days of exposure fragmentation of some red blood cells was clear (Fig. 1b), with appearance of tear drop like cells and acanthocytes. After 12 and 16 days of exposure the red blood cells lose their normal oval shape so that they become nearly rounded. Crenated cells acanthocytes and tear drop like cells were frequent (Fig. 1c). Beside lysis of some red blood cells, some cells appear to be stacked together.

(2) Effect of sublethal concentration of Cadmium (6.5 ppm):

In the blood film of fish exposed to sublethal concentration of Cd (6.5 ppm) for 4 days it appears that most of the red blood cells lose their normal oval shape (Fig. 2a). Acanthocytes and crenated cells were apparent beside lysis of some red blood cells. Sticking of the red blood cells and fragmentation of red blood cells were also observed. Upon further exposure (8 days) the phenomena of sticking of the cells is further apparent (Fig. 2b). Acanthocytes, Crenated cells, tear drop cells and sickle cells were apparent. After 12 days of exposure sticking of the cells become more apparent. The stacked cells lose their original shape with appearance of sickle shaped cells (Fig. 2c). The nuclei of the red blood cells shows some variation in their shape. By 16 days of exposure cells acquired different shapes with obvious disintegration of cell wall of some cells and acanthocytes were frequent (Fig. 3a).

(3) Effect of exposure to sublethal concentration of copper (0.7 ppm):





The first observed change in the red blood cell morphology after treatment with Cu was the rouleaux appearance that appeared in the first 4 days of treatment Fig. 3b). Also cells lose their normal shape acquiring different shapes or become rounded (Fig. 3c). After 8 days of exposure the red blood cells started to be stacked with each other with nuclei losing some amount of central dye showing a sign of karyolysis (Fig. 4a) and significantly increase in size ($t = 3.5$ and $P < 0.5$). Sick cells were apparent and most of erythrocyte cells lose their normal shape. Upon 12 days of exposure most of the red blood

cells lose their normal oval shape and become polygonal or irregular in shape (Fig. 4b). Sick cells were apparent. Also spherocyte and tear drop like cells were observed (Fig. 4c).

(II) Variations in R.B.Cs count and haemoglobin content:

(1) Zinc

R.B.Cs count significantly decreased than that of control fish after 4, 8 and 12 days of exposure to sublethal concentration of Zinc (4.6 ppm) ($t = -5.8$ (4 d), -25.7 (8 d) & -13.4 (12 d) $P < 0.05$). A significant decrease in haemoglobin content as compared to control fish also occurred ($t = -9.25$ (4 d), -10.66 (8 d) and -16.48 (12 d) $p < 0.05$).






Figure shows the effect of Sublethal concentration of Zn, Cd and Cu on normal blood cells.

(2) Cadmium

Significant decrease in RBCs count was noticed ($t = -3.9$ (4 d), -14 (8 d) and -3.6 (12 d) $P < 0.05$). A significant decrease in hemoglobin content hemoglobin content of fish also occurred ($t = -14.9$ (4 d), -23.1 (8 d) and -16.5 (12 d) $P < 0.05$).

(3) Copper

A slight increase in RBCs of the fish that exposed to 0.7 ppm Cu after 4, 8 and 12 day was observed ($t = -4.07$ (4 d), -4.2 (8 d) and -3.05 (12 d) $P < 0.05$). Significant increase in haemoglobin content was also noticed ($t = 8.2$ $P < 0.05$) after 4 days of exposure. However this rise did not continue, either after 8 days or 12 days.

4. Discussion

The effect of pollution on the blood profile of fishes made the significant studies of various workers (Osman *et al.*, 1993 ^[25], Zia and McDonald 1994 and Adak, 1995 ^[5]). Comparison of haematological characteristics of *C. batrachus* with those of other fishes are given according to the present study exposure of *C. batrachus* to sublethal concentrations of heavy metals Zn (2.3 ppm), Cd (8.4 ppm) and Cu (0.2 ppm) resulted in some

changes in the morphology of the R.B.Cs which tended to lose its ellipsoidal shape acquiring different shapes. Among the abnormal cells were the tear drop like cells, acanthocytes, crenated cells and sickle cells. Another effects of metals observed in the present study, are the sticking of the cells together, decrease (in case of Zn) or increase (in case of Cu) in size and appearance of spherocytes. In case of zinc and cadmium cell wall of the erythrocytes may disintegrate leaving the nuclei free in the blood film, as well as cell fragments. Few studies are concerned with study of the morphological changes in red blood cells of fishes upon exposure to heavy metals or pollutants as a whole according to available literature. Smith (1968) [28] recorded poikilocytosis and anisocytosis of red blood cells of coho salmon *Onchorhynchus kistuch* fed on folic acid deficient diet. He claimed that these results are characteristic feature of megaloblastic, normoblastic and macroblastic anaemia. Srivastava and Mishra (1978) found that exposure of *Colisa fasciatus* to sublethal concentration of lead resulted in lysis of the erythrocytes. Ahmed and Munshi (1992) [5] have found that exposure of Indian carp *Catta catta* to copper resulted in shrinkage and crenation of the configuration of the red blood cells with slight anisocytosis and with a tendency to overlap. Poikilocytosis of red blood cells resulted in association with various types of anaemia e.g. megaloblastic anaemia, iron deficiency anaemia, thalassemia, myeloclerosis or from damage to circulating red blood cells as in microangiopathic hemolytic anaemia (Dacie and Lewis, 1975) [11]. This suggests that fish exposed to heavy metals suffer from anaemia.

Acanthocytosis is seen in the liver diseases (Cooper, 1980) [10] or in association with abnormal phospholipid metabolism (Dacie and Lewis, 1975) [11]. This suggests that the fishes exposed to heavy metals (at sublethal doses) suffer from liver disease. Tear drop cells may result from the Pitting action of the spleen which removes autophagocytic vacuoles or inclusions. They are also seen in beta thalassemia, Heinz body anaemia, megaloblastic processes and disorders or leukemia and are caused by the removal of precipitated haemoglobin inclusions by the spleen (Bike, 1993) [8]. Sick cells are present in homozygous Hb-S disease in films of freshly withdrawn blood and subjected to anoxia (Dacie and Lewis, 1975) [11].

Exposure of *C. batrachus* to sublethal concentrations of Zn and Cd resulted in significant decrease ($p < 0.05$) in erythrocyte count and haemoglobin content. This case was parallel to the observation of Smith *et al.*, (1971) on chinook salmon *Onchorhynchus tshawytscha* subsequent to treatment with phenylehydrazin; in coho salmon *Onchorhynchus kistuch* following exposure to residual chlorine (Buckly *et al.*, 1976) and generally in teleosts exposed to pulp mill effluent (Warner, 1967 [30] and McLeay, 1975 [24]) and to cadmium (Larsson *et al.*, 1976). The same observation was also recorded on *Colisa fasciatus* following exposure to lead (Srivastava and Mishra, 1979) [26], in *Clarias lazera* after acute exposure to lead (Shabana, 1983) on *Anguilla vulgaris* after exposure to mercury (Hilmy *et al.*, 1987) [18], on *Clarias lazera* exposed to hexavalent chromium (Hilmy *et al.*, 1988) [19] on *Oreochromis niloticus* exposed to cadmium (AL-Akel *et al.*, 1988) [6], on *Carassius carassius* exposed to lead (Fantin *et al.*, 1989) [13], on *Clarias lazera* exposed to mercury and gallant (Abd-Allah *et al.*, 1991 [1]) on European silver eel exposed to lead (Amin, 1992) [7] and on *Oreochromis mossambicus* exposed to arsenic and zinc (Adak, 1995) [5].

Abdel-Maguid (1997) have found that starvation resulted in

decrease of both R.B.Cs count and haemoglobin content of *Gobius niger*. On the other hand upon exposure of *C. batrachus* to sublethal concentration of copper significant increase in erythrocyte count and haemoglobin content was observed. Ghazaly and Said (1995) [16] have found similar results in *Tilapia nilotica* after exposure to sublethal concentration of copper. Some other metals were found to induce the same effect of copper on fish blood. Thus Frovola (1960) [14] had recorded that an increase in erythrocyte count and haemoglobin content in carp *Cyprinus carpio* after exposure to cobalt. Similar type of results were found in rainbow trout *Salmo gairdneri* (Schiffman and Forman, 1959 [27]) and in *Colisa fasciatus* (Srivastava *et al.*, 1979) following exposure to chromium.

5. Conclusion

Exposure of *C. batrachus* to sublethal concentrations of Zn and Cd resulted in significant decrease ($p < 0.05$) in erythrocyte count and haemoglobin content, while there was slight increase in case of copper. The microscopical observations showed that exposure of *C. batrachus* to sublethal concentrations of heavy metals Zn (3.8 ppm), Cd (7.6 ppm) and Cu (0.4 ppm) resulted in some changes in the morphology of the R.B.Cs which tended to lose its ellipsoidal shape acquiring different shapes (poikilocytosis or anisocytosis). Among the abnormal cells were the tear drop like cells, acanthocytes, crenated cells and sickle cells. Another effects of metals observed in the present study, are the sticking of the cells together, decrease (in case of Zn) or increase (in case of Cu) in size and appearance of spherocytes. In case of copper and cadmium cell wall of the erythrocytes may disintegrate leaving the nuclei free in the blood film, and appearance of cell fragments.

6. Acknowledgement

Authors are beholden to Principal Dr. A.K. Gangely S.L.L. Jain P.G. College Vidisha, Bhopal M.P. India for providing laboratory facilities for this research work. This paper forms the part of Ph.D. thesis submitted by author to Barkatullah University, Bhopal M.P. (India).

7. References

- 1 Abd-Allah GA, Ibrahim MS, Bahnasawy MH, Abdel-Baky TF. Toxic effects of some water pollutants (Gallant and mercury) on blood parameters of cat fish *Clarias Lazera*. J Egypt Ger Soc Cool. 1991; 6(A):201-209.
- 2 Abdel-Barr M. Toxicological studies of heavy metals on *Clarias batrachus*. M. Sc. thesis, Fac. Sci. Alex. Univ. 1997, 188.
- 3 Abdel-Maguid SA. Biological and haematological studies on marine coastal fish, family, Gobiidae in Alexandria, Egypt. Ph.D. Thesis, Fac. Sci. Alex. Univ. 1997, 206.
- 4 Adak SK. Haematological studies on fresh water telost, *Oreochromis (Tilapia) mosambicus* (Peters) in the normal conditions and under the stress of zinc and Arsenic. Ph.D. Thesis, zoology, Sambalpur Univ. 1995, 64.
- 5 Ahmed IM, Munshi S. Scanning electron microscopic evaluation of changes on the morphology of blood cells of an Indian major carp *Catta catta* (Halm). Following exposure to copper. J Environ Biol. 1992; 13(14):297-301.
- 6 Al-Akel AS, Shami MJ, Al-Kalem HF, Ghaudhary MA, Ahmed Z. Effects of cadmium on the cichlid fish *Oreochromis niloticus* behavioural and physiological responses. J Univ Kuwait Sci. 1988; 5(2):341-345.

- 7 Amin EM. Hematological changes and light microscopic study of peripheral blood cells in European eel exposed to lead concentrations. Rapp. Comm. nt. Mer. Medit. 1992; 33:165.
- 8 Bike RL. Hematology: Clinical and Laboratory practice 1:12 Morphology of erythron, St.Louis, Missouri, 1993, 446.
- 9 Buckley JA, Whitenore CM, Matsuda RI. Changes in blood chemistry and blood cell morphology in Coho-Salmon *Oncorhynchus Kisutch* following exposure to sublethal levels of total residual chlorine in municipal waste water. J Fish Res Can. 1967; 33:776-782.
- 10 Cooper RA. Red cell membrane abnormalities in liver disease. Semin, Hematol. 1980; 17:103.
- 11 Dacie JV, Lweis SM. Practical hematology. 5th ed. Churchill Livingstone, 1975, 564.
- 12 Dawson AB. The hemopoietic response in the cat fish *Ameiurus nebulosus*, to chronic lead poisoning. Biol. Bull. 1953; 68:335-346.
- 13 Fantin AMB, Tervisan P, Pederzoli A, Bergoni M. Effect of experimental pollution by lead on some hematological parameters in *Carassius carassius* var *anratus*. Bollctino Dizologia. 1989; 55(4):251-256.
- 14 Frovola LK. The effect of inorganic cobalt on the morphological picture of *Cyprinus carpio*. Dolel. Akad, Nauk. SSSR. 1960; 131:933-984.
- 15 Gardner GR, Yevich PP. Toxicological effects of cadmium on *Fundulus heteroclitus* under various oxygen, Ph., Salinity and temperature regimes. Amer. Zool. 1969b, 1096.
- 16 Ghazaly KS, Said KM. Physiological characteristics of *Tilapia nilotica* under stress of copper. J Egypt Ger Soc. Zool. 1995; 54(3):78-81.
- 17 Hesser EF. Methods of routine fish hematology. Progressive fish culturist, 1960; 22(4):164-171.
- 18 Hilmy AM, Shabaua MB, El-Domiaty, NA, Yaacout LE. Mercury levels and hematological correlation in *Anguilla Vulgaris*. Bull. Fac. Sci. Alex. Univ. 1987; 27(4), A:125-140.
- 19 Hilmy AM, El-Domiaty MA, Moussa FI, El-Masry AA. Hematological and physical changes induced by Hexavalent chromium in *Clarias Lazera*. Bull. Inst. Oceanogr. & Fish. A.R.E. 1988; 14(3):101-110.
- 20 Houston H, Korm MD, Cohen Y, Bernhard M. Trace metal content in deep water sharks in eastern Mediterranean sea Marin. Biol. 1993; 115:31-338.
- 21 Khardre SM. Changes in the gill structure and blood profiles following acute copper toxicity in two fresh water teleosts. Proc. int. 1990; 44(4):27-31.
- 22 Larrson A, Bengtsson BE, Svanbery A. Some hematological and biochemical effects of cadmium on fish. From the effect of pollutants on aquatic organisms. A:P.M. Lockwood. Cambridge University. Press, 1976.
- 23 McCay CM. Studies upon fish blood and its relation to water pollution. A Biological survey for the Erie-Niagra system. New York conservation Dep., N.Y. 1929, 140-149.
- 24 McLeay DJ. Sensitivity of blood cell count in juvenile coho salmon *Onchorhynchus Kistuch*. J Fish Res Bd Can. 1975; 32:2357-2364.
- 25 Osman MA, Ghaly YS, Afify AMR, El-Dosoky G, Mahmoud MEF. Lend pollution of water resources and its biochemical effects. Proc. 3rd. Int. Conf. on Environ. Prot. 1993, 75-87.
- 26 Sarivastava AK, Mishra S. Blood dyscrasia in teleost, *Colisa fasciatus* after acute exposure to sublethal concentration of lead. J. Fish Biol. 1979; 14:199-203.
- 27 Schiffman RH, Forman PO. Chromium induced changes in the blood of rainbow trout *Salmo gairdneri*. Sewage and industrial wastes. 1959; 31: 205-211.
- 28 Smith C. Hematological changes in cho salmon fed on folic acid deficient diet. J. Fish Res. Bd. Can. 1968; 25(1):151-156.
- 29 Srivastava AK, Agrawal SJ, Ghaudhry HS. Effects of chromium on blood of fresh water fish. Ecotoxicol. Environ. saf. 1979; 3:321-324.
- 30 Warner RE. Bioassay for micro chemical environments. Bull. W.H.O. 1967, 181-207.
- 31 Wintrow MM. Clinical hematology. London, Henry Kimpton. 1967, 448.