



International Journal of Fisheries and Aquatic Studies

E-ISSN: 2347-5129

P-ISSN: 2394-0506

(ICV-Poland) Impact Value: 5.62

(GIF) Impact Factor: 0.549

IJFAS 2018; 6(3): 93-95

© 2018 IJFAS

www.fisheriesjournal.com

Received: 12-03-2018

Accepted: 13-04-2018

Periklis Kleitou

(1) iSea, Environmental
Organization for the
Preservation of the Aquatic
Ecosystems, Greece

(2) Marine and Environmental
Research (MER) Lab Ltd.
Limassol, Cyprus

(3) School of Marine Science and
Engineering, Plymouth
University, Plymouth, UK

Stefanos Kalogirou

Hellenic Centre for Marine
Research, Hydro biological
Station of Rhodes, 85100
Rhodes, Greece

Dimitra Marmara

Marine and Environmental
Research (iSea) Lab Ltd.
Limassol, Cyprus

Ioannis Giovos

Marine and Environmental
Research (iSea) Lab Ltd.
Limassol, Cyprus

Correspondence

Periklis Kleitou

(1) iSea, Environmental
Organization for the
Preservation of the Aquatic
Ecosystems, Greece

(2) Marine and Environmental
Research (MER) Lab Ltd.
Limassol, Cyprus

(3) School of Marine Science and
Engineering, Plymouth
University, Plymouth, UK

Coryphaena hippurus: A potential predator of *Lagocephalus sceleratus* in the Mediterranean Sea

Periklis Kleitou, Stefanos Kalogirou, Dimitra Marmara and Ioannis Giovos

Abstract

Lagocephalus sceleratus is considered as one of the most invasive Lessepsian immigrants introduced in the Mediterranean Sea. It is a highly toxic fish, considered as a pest for fisheries, the environment and the economy. Hitherto, its invasion has been favoured by the lack of natural predators and its non-commercial value. The present study reports the first published incidence of a juvenile *L. sceleratus* predation in the Mediterranean by an adult *Coryphaena hippurus*. This constitute a significant finding which can potentially guide future management efforts of *L. sceleratus* control in the basin; however, further research should be conducted to elucidate whether the catch was accidental or if the sustainable management of *C. hippurus* populations can potentially affect the early life stages of *L. sceleratus*.

Keywords: Mediterranean, lagocephalus, predation, mahi mahi, invasive, management

1. Introduction

Lagocephalus sceleratus (Gmelin, 1789), known as the silver-cheeked toadfish, is widely distributed widely in the tropical and subtropical waters of the Indian and Pacific Ocean, including the southern African shores and the Red Sea, recently entered in the Mediterranean Sea. Since its first record in the basin in 2003^[1], this Lessepsian immigrant has established populations along the coasts of the eastern Mediterranean basin, and expanded towards the western Basin; being recorded from Spain in 2014^[2,3].

The species has received considerable attention from the public and the scientific community as it contains high concentrations of tetrodotoxin (TTX) in its tissues, which can be fatal when consumed^[4,5]. Although regulations have been imposed to prohibit sale and consumption of *Tetraodontidae*, such as *L. sceleratus* (e.g. EC 854/2004), more than 15 incidences of human intoxication have been reported from the eastern Mediterranean countries so far, most recently from Cyprus and Greece^[6,7]. *L. sceleratus* is also considered to cause ecological and economic damages and viewed as a pest by fishermen; capable of reducing the local stocks of important commercial cephalopod species, damaging fishing gears, deterring customers from buying fish and introducing additional effort to discard the fish^[8-10]. These impacts have led scientists to classify *L. sceleratus* invasion among the 100 worst marine invasions in the Mediterranean basin^[11].

The non-commercial value of *L. sceleratus* and possible lack of natural predators^[12] may have contributed to its rapid expansion in the Mediterranean Sea. To this end, some countries, such as Cyprus, have even provided financial compensation (i.e. €1-3 per kg) to encourage fishermen to target the fish in an effort to control its population^[13]. To our knowledge, there is no previous documented information on natural predators for the species in the Mediterranean basin. Here, the first published incidence of *L. sceleratus* predation in the Mediterranean Sea is reported.

2. Materials and methods

In 2016, an online data repository was established by iSea (i.e. "Is it Alien to you.... Share it!!!"), in which citizen scientists could easily upload photographic material along with information on specimen size (length and/or weight), depth, number of specimens, exact location, date and type of observation (freediving, underwater photography, shore-base fishing, boat-based fishing, spearfishing). A Google Form and a Group on Facebook were established to facilitate these reports.

3. Results and Discussions

On 29 August 2017, a fisherman reported an incidence of potential predation of *L. sceleratus* through the citizen-science programme. Specifically, one individual of *Coryphaena hippurus* Linnaeus, 1758 was captured at 09:50 a.m. using shore angling, at Plakias harbour, South Rethymno (Crete, Greece) (35°11'34.4"N; 24°22'51.7"E) (Figure 1). The fish was found swimming in a school and caught over a sandy substrate at 4 m depth. The specimen weighted around 2-3 kg and was approximately 70 cm (fork) length (Figure 2A; B). A juvenile *L. sceleratus* of approximately 3 cm was found in the oesophagus. The prey appeared freshly consumed with no signs of degradation or digestion (Figure 2C; D). All the information and evidences were submitted to the programme.




Fig 1: The catch location of the *C. hippurus* that consumed a juvenile *L. Sceleratus* in South Crete, Greece.



Fig 2: The specimen of *C. hippurus* caught (A, B) and the juvenile *L. sceleratus* (C, D) found in the oesophagus.

It has been demonstrated that non-toxic species show low resistance to TTX and TTX-bearing organisms use it effectively as a defensive or offensive substance [14]. According to the results of Katikou *et al.* (2009) [15], toxicity was not detected in any of the tested tissues of two *L. sceleratus* specimens smaller than 16 cm. However, the effect of *L. sceleratus* consumption on other species acting as predators has not been extensively studied and therefore, any assumption would be ambiguous.

Studies in the Pacific and Atlantic Oceans and the Arabian Sea have found that *C. hippurus* may consume a variety of Tetraodontidae species, including the congeneric toxic species *L. lagocephalus* (Linnaeus, 1758) [16-20]. An earlier study on the diet of *C. hippurus* in the western Mediterranean Sea found a large proportion of epipelagic prey and indicated that

C. hippurus is a top predator, but no species of Tetraodontidae were reported [21]. However, many Tetraodontidae species such as *L. sceleratus* have been expanded into the Mediterranean after the study of [21] and it is therefore imperative that updated studies on the diet of *C. hippurus* are being conducted.

4. Conclusions

Studies on the diet of *C. hippurus* could usefully be undertaken to determine whether the species is a potential natural predator of juvenile *L. sceleratus* in the Mediterranean. Such information can be vital for managers as it can guide future strategies towards the control and management of the invasive *L. sceleratus* in the basin.

5. Acknowledgments

The authors are grateful to the fisherman Mr Konstantinos Tsagkarakis (Greece), for sharing his information and pictures to the citizen-science programme "Is it Alien to you.... Share it."

References

1. Filiz H, Er M. Akdenizin yeni misafiri (New guests in the Mediterranean Sea). *Deniz Magazin (Istanbul)*. 2004; 68:52-4.
2. Katsanevakis S, Acar Ü, Ammar I, Balci B, Bekas P, Belmonte M *et al.* New mediterranean biodiversity records (October, 2014). *Mediterranean Marine Science*. 2014; 15(3):675-95.
3. Deidun A, Fenech-Farrugia A, Castriota L, Falautano M, Azzurro E, Andaloro F. First record of the silver-cheeked toadfish *Lagocephalus sceleratus* (Gmelin, 1789) from Malta. *BioInvasions Records*. 2015; 4(2):139-42.
4. Kosker AR, Özogul F, Durmus M, Ucar Y, Ayas D, Regenstein JM *et al.* Tetrodotoxin levels in pufferfish (*Lagocephalus sceleratus*) caught in the Northeastern Mediterranean Sea. *Food chemistry*. 2016; 210:332-7.
5. Rambla-Alegre M, Reverté L, Del Río V, de la Iglesia P, Palacios O, Flores C *et al.* Evaluation of tetrodotoxins in puffer fish caught along the Mediterranean coast of Spain. *Toxin profile of Lagocephalus sceleratus*. *Environmental research*. 2017; 158:1-6.
6. Bentur Y, Ashkar J, Lurie Y, Levy Y, Azzam ZS, Litmanovich M *et al.* Lessepsian migration and tetrodotoxin poisoning due to *Lagocephalus sceleratus* in the eastern Mediterranean. *Toxicon*. 2008; 52(8):964-8.
7. Katikou P, Vlamis A. Tetrodotoxins: recent advances in analysis methods and prevalence in European waters. *Current Opinion in Food Science*. 2017; 18:1-6.
8. Kalogirou S, Corsini-Foka M, Sioulas A, Wennhage H, Pihl L. Diversity, structure and function of fish assemblages associated with *Posidonia oceanica* beds in an area of the eastern Mediterranean Sea and the role of non-indigenous species. *Journal of fish biology*. 2010; 77(10):2338-57.
9. Kalogirou S, Wennhage H, Pihl L. Non-indigenous species in Mediterranean fish assemblages: Contrasting feeding guilds of *Posidonia oceanica* meadows and sandy habitats. *Estuarine, Coastal and Shelf Science*. 2012; 96:209-18.
10. Kalogirou S. Ecological characteristics of the invasive pufferfish *Lagocephalus sceleratus* (Gmelin, 1789) in the eastern Mediterranean Sea—a case study from Rhodes. *Mediterranean Marine Science*. 2013; 14(2):251-60.

11. Streftaris N, Zenetos A. Alien marine species in the Mediterranean-the 100 'Worst Invasives' and their impact. *Mediterranean Marine Science*. 2006; 7(1):87-118.
12. East MED F. Report of the sub-regional technical meeting the lessepsian migration and its impact on eastern Mediterranean fishery. Nicosia, Cyprus, December, 2010, 5-7.
13. Rousou M, Ganias K, Kletou D, Loucaides A, Tsinganis M. Maturity of the pufferfish *Lagocephalus sceleratus* in the southeastern Mediterranean Sea. Sexuality and early development in Aquatic organisms. 2014; 1(1):35-44.
14. Nagashima Y, Arakawa O. Pufferfish Poisoning and Tetrodotoxin. *Marine and Freshwater Toxins*, 2016, 259-84.
15. Katikou P, Georgantelis D, Sinouris N, Petsi A, Fotaras T. First report on toxicity assessment of the Lessepsian migrant pufferfish *Lagocephalus sceleratus* (Gmelin, 1789) from European waters (Aegean Sea, Greece). *Toxicon*. 2009; 54(1):50-5.
16. Olson RJ, Galván-Magaña F. Food habits and consumption rates of common dolphinfish (*Coryphaena hippurus*) in the eastern Pacific Ocean. *Fishery Bulletin*. 2002; 100(2):279-98.
17. Tripp-Valdez A, Galvan-Magaña F, Ortega-Garcia S. Food sources of common dolphinfish (*Coryphaena hippurus*) based on stomach content and stable isotopes analyses. *Journal of the Marine Biological Association of the United Kingdom*. 2015; 95(3):579-91.
18. Varela JL, Lucas-Pilozo CR, González-Duarte MM. Diet of common dolphinfish (*Coryphaena hippurus*) in the Pacific coast of Ecuador. *Journal of the Marine Biological Association of the United Kingdom*. 2017; 97(1):207-13.
19. Varghese S, Somvanshi V, John M, Dalvi R. Diet and consumption rates of common dolphinfish, *Coryphaena hippurus*, in the eastern Arabian Sea. *Journal of Applied Ichthyology*. 2013; 29(5):1022-9.
20. Brewton RA, Ajemian MJ, Young PC, Stunz GW. Feeding Ecology of Dolphinfish in the Western Gulf of Mexico. *Transactions of the American Fisheries Society*. 2016; 145(4):839-53.
21. Massutí E, Deudero S, Sánchez P, Morales-Nin B. Diet and feeding of dolphin (*Coryphaena hippurus*) in western Mediterranean waters. *Bulletin of Marine Science*. 1998; 63(2):329-41.