



# International Journal of Fisheries and Aquatic Studies

ISSN: 2347-5129

IJFAS 2015; 2(4): 299-303

© 2015 IJFAS

www.fisheriesjournal.com

Received: 18-02-2015

Accepted: 02-03-2015

## Deniz Innal

Mehmet Akif Ersoy University,  
Department of Biology, Istiklal  
Campus. 15100, Burdur Turkey.

## Mehmet Aksu

Mehmet Akif Ersoy University,  
Department of Biology, Istiklal  
Campus. 15100, Burdur Turkey.

## Duygu Akdoganbulut

Mehmet Akif Ersoy University,  
Department of Biology, Istiklal  
Campus. 15100, Burdur Turkey.

## Burcu Kisin

Mehmet Akif Ersoy University,  
Department of Biology, Istiklal  
Campus. 15100, Burdur Turkey.

## Mehmet Can Unal

Mehmet Akif Ersoy University,  
Department of Biology, Istiklal  
Campus. 15100, Burdur Turkey.

## Mustafa Oztop

Mehmet Akif Ersoy University,  
Department of Biology, Istiklal  
Campus. 15100, Burdur Turkey.

## Bugrahan Dogangil

Mehmet Akif Ersoy University,  
Department of Biology, Istiklal  
Campus. 15100, Burdur Turkey.

## Engin Pek

Mehmet Akif Ersoy University,  
Department of Biology, Istiklal  
Campus. 15100, Burdur Turkey.

## Correspondence

### Deniz Innal

Mehmet Akif Ersoy University,  
Department of Biology, Istiklal  
Campus. 15100, Burdur Turkey.

## Age and growth of *Nemipterus randalli* from Antalya Gulf-Turkey

**Deniz Innal, Mehmet Aksu, Duygu Akdoganbulut, Burcu Kisin, Mehmet Can Unal, Mustafa Oztop, Bugrahan Dogangil, Engin Pek**

### Abstract

Age, growth, length-length and length-weight relationships were estimated for *Nemipterus randalli* Russell, 1986 specimens captured between September 2012 and April 2014 from Gulf of Antalya, Turkey. We examined a total of 175 individuals, of which range in size between 6 and 24 cm in total length, 3.4 and 201 g in total weight. Maximum ages were III for female and IV for male. The overall male to female ratio was 0.90/1.0. The length-weight relationship for all individuals were described by the parameters:  $a = 0.0105$  and  $b = 3.0426$ . Von Bertalanffy growth equations were estimated as  $L_t = 33.15[1 - e^{-0.218(t+0.215)}]$  and  $W_t = 444[1 - e^{-0.218(t+0.215)}]^{3.0426}$  and the phi-prime test estimation was calculated as 2.38 in the population. This study also presents the relationships between total length-standard length, total length-fork length, standard length-fork length, total length-head length, total length-head depth, total length-body depth, total length-otolith length, total length-otolith width, body depth-head depth, head length-head depth and body depth-head length for *N. randalli*.

**Keywords:** *Nemipterus randalli*, age, growth, length-length and length-weight relationships, Antalya Gulf

### 1. Introduction

Non-indigenous fish species have become established in various parts of the inland waters and coasts of Turkey. The colonization of Red Sea species in the Mediterranean Sea is an ongoing process that began some time after the opening of the Suez Canal in 1869. *Nemipterus randalli* is one of the Lessepsian migrant species, first recorded by Golani and Sonin (2006) [14] on the Mediterranean coast of Israel mistakenly as *Nemipterus japonicus*. However, *Nemipterus japonicus* have nowadays been recognized as an invasive species from the Mediterranean [7], together with *N. randalli*.

Tropical and subtropical regions are the ideal habitat for Nemipterid fishes and this family (Nemipteridae) includes 67 species belonging to 5 genera, namely *Nemipterus*, *Parascolopsis*, *Pentapodus*, *Scaevius*, and *Scolopsis* [12]. The threadfin breams of the genus *Nemipterus* (family Nemipteridae) are widespread species that occur at Indo-West Pacific region, including the western Indian Ocean, the east and west coasts of India, the Persian Gulf, and the Red Sea south to Madagascar and more than 20 species are recognized [27]. They are small to moderate-sized fishes and most inhabit shallow sand or mud bottoms, and are taken commercially by hook-and-line and bottom trawl [28].

Various biological parameters of *N. randalli* have been studied in the Mediterranean by various authors; for example, distribution [14, 20, 5, 13, 1, 15] diet [16], population structures and some growth properties [20, 5, 13, 16, 15, 1, 2, 23].

*Nemipterus* species are commercially important in many parts of the world [9]. This species has rapidly extended and now successfully established in Eastern Mediterranean Sea [16]. There is no catch statistics for threadfin bream in Antalya Gulf Turkey, although the abundance of this species rapidly increase (pers observation, D. Innal).

Although *N. randalli* has been studied along the Mediterranean Sea, there has been no previous references for biological properties from Antalya Gulf. The objective of this study was to provide information on the length-length, length-weight relationships and some growth properties of *N. randalli* in the Antalya Gulf.

## 2. Material and Methods

### 2.1 Data collection and analysis

Samples (175 individuals) were collected during September 2012 and April 2014 from trawl surveys conducted in the Antalya Gulf (Mediterranean-Turkey). Fish samples were immediately transported to the laboratory in the Department of Biology, Mehmet Akif Ersoy University (Turkey). Size (total length cm TL, precision 1 mm) and weight (total weight W to nearest 0.1 g) were calculated. Sex was identified macroscopically and microscopically by examining the gonads. Metric characters were measured with digital slide calliper on the fish body (4 characteristics; total length, standard length, fork length, body depth at dorsal fin), head (4 characteristics; head length, head depth, otolith length, otolith width). Morphometric relationships between parameters (total length-standard length, total length-fork length, standard length-fork length, total length-head length, total length-head depth, total length-body depth, total length-otolith length, total length-otolith width, body depth-head depth, head length-head depth and body depth-head length) were calculated using the linear regression.

For age determination, 175 sagittal otoliths were extracted, washed, dried, and mounted on black slides. A stereo microscope with reflected light was used for age determination. Length-weight relationship was calculated using the equation  $W = aL^b$  [24], where  $a$  is a coefficient related to body form and  $b$  is an exponent indicating isometric growth when equal to 3. The  $b$ -value of each species was tested by Student's  $t$ -test to verify if it was significantly different from isometric growth [11]. The von Bertalanffy growth equations were calculated according to:  $L_t = L_\infty[1 - e^{-k(t-t_0)}]$  in length and  $W_t = W_\infty[1 - e^{-k(t-t_0)}]^b$  in weight, where  $L_t$  is the fish total length (cm) at age  $t$ , 'e' is the base of natural log (2.71828), 't' the fish age (in years) to the hypothetical time at which the length of the fish is zero, and  $k$  is a relative growth coefficient.  $W_t$  is the fish weight (g) at age  $t$  [24]. Munro's phi prime index (growth performance index) was estimated by formula  $\Phi' = \log(k) + 2\log(L_\infty)$  [25]. Differences in Size frequency distributions for males and females were calculated by  $t$ -test. Comparisons if sex ratio departed from the expected 1 : 1 rate were determined by  $t$ -test.

## 3. Results

### 3.1 The Sex and Age Compositions

The sex and age compositions of *N. randalli* are shown in Table 1. The 175 *N. randalli* individuals consisted of 76 (43.43%) males, 84 (48%) females and 15 (8.57%) immature individuals, sex ratio is significantly different ( $P < 0.05$ ) from the expected 1 : 1 (male, 0.90: female, 1). Ages of captured specimens ranged from 0 to IV, with first year-class being dominant in the population.

### 3.2 Length and Weight Composition

Length and weight distribution and Length-weight relationships (LWR) of *N. randalli* are given in Fig. 1. Age-length and age-weight composition of *N. randalli* are shown in Table 1. Specimens of *N. randalli* ranged from 6 to 24 cm TL and from

3.4 to 201 g weight. Among all individuals, dominance (66.8%) was in the size range of 13–17 cm. Adult males ranged from 12 to 24 cm TL and females from 12.2 to 22.9 cm TL. Size frequency distributions for males and females were not significantly different ( $t$ -test,  $P > 0.05$ ). The length-weight relationship for all individuals were described by the parameters:  $a = 0.0105$  and  $b = 3.0426$ . LWRs were also calculated for sexes separately. In *N. randalli* these were  $W = 0.008 L^{3.1365}$  ( $R^2 = 0.97$ ) for females and  $W = 0.0079 L^{3.1498}$  ( $R^2 = 0.98$ ) for males.




Fig 1: Length and weight distribution and Length-weight relationships (LWR) of *N. randalli*

**Table 1:** Sex and age compositions; Age-length and Age-weight key of *N. randalli*

Length classes	age classes					N
	0	1	2	3	4	
5-6.9	1					1
7-8.9	1					1
9-10.9	7	4				11
11-12.9		12				12
13-14.9		60				60
15-16.9		18	39			57
17-18.9			6	1		7
19-20.9			14	4		18
21-22.9			1	2	3	6
23-24.9					2	2
5-24.9	9	94	60	7	5	175
Sex ratio (f/m)	immature	0.80/1	2.33/1	0.75/1	0/1	1.11/1
Length						
Mean±std	8.92±1.41	13.90±1.22	17.26±1.59	20.39±1.27	23.06±0.85	14.9±2.44
Min	6	10.3	15.7	18.9	21.9	9
Max	10.1	15.8	21.5	22.9	24	24
Weight						
Mean±std	8.63±3.06	31.93±7.42	62.79±22.19	108.6±27.9	165.8±25.14	42.22±26.24
Min	3.4	14.2	41.8	77.2	134	7.7
Max	12.5	47.5	130	160	201	201

### 3.3 Length- length relationships

Regression analysis between the different body and head measurements for *N. randalli* are shown in Table 2. All Length-

length relationships were highly significant with  $r^2$  values greater than 0.87. Except only one relationships was found with  $r^2= 0.74$  (otolith length-total length relationships)

**Table 2:** Length- length relationships and correlation coefficients for *N. randalli* from the Antalya Gulf (Mediterranean-Turkey).

Correlates	Equations	$r^2$
TOTAL LENGTH - FORK LENGTH	TL = 1.066 (FL) +0.9032	0.98
TOTAL LENGTH - STANDART LENGTH	TL = 1.1026 (SL) +2.1023	0.94
FORK LENGTH - STANDART LENGTH	FL = 1.0051 (SL) +1.4643	0.93
TOTAL LENGTH - BODY DEPTH	BD = 0.2405 (TL) -0.224	0.87
TOTAL LENGTH - HEAD LENGTH	HL = 0.2722 (TL) -0.4212	0.93
TOTAL LENGTH - HEAD DEPTH	HD = 0.2412 (TL) -0.8785	0.91
BODY DEPTH - HEAD DEPTH	HD = 0.833 (BD) -0.0827	0.87
HEAD LENGTH - HEAD DEPTH	HD = 0.8436 (HL) -0.3498	0.93
BODY DEPTH - HEAD LENGTH	HL = 0.9175 (BD) +0.5521	0.87
OTOLITH LENGTH - TOTAL LENGTH	OL = 0.3486 (TL) +0.7416	0.74
OTOLITH WIDTH - TOTAL LENGTH	OW = 0.2404 (TL) +0.4728	0.87

### 3.4 Von Bertalanffy growth

The von Bertalanffy growth model of the *N. randalli* population in Antalya Gulf was described as  $L_t = 33.15[1 - e^{-0.218(t+0.215)}]$  and  $W_t = 444[1 - e^{-0.218(t+0.215)}]^{3.0426}$ . The phi-prime test estimation was calculated as 2.38 in the population.

### 4. Discussion

During the present study females were more abundant in the catch of *N. randalli*. The sex ratio, which differs from one population to another in the same species, was 1.0 females to 0.90 male in the *N. randalli* populations in present study. The sex ratio (female:male) in previous studies of *N. randalli* populations was as follows: Iskenderun Bay, 1.29:1<sup>[9]</sup>; Arabian Sea coast of Oman, 0.9:1<sup>[2]</sup>. The sex ratio may vary from year to year in the same population, indicating that it is either determined genetically or by environmental factors<sup>[4]</sup>. This variation may arise from many reasons, including seasonal aspect, feeding and maturation periods, different growth rates in males and females, mortality difference for each sex, and perhaps, size-selective effects of the fishing gear.

The life span of males and females differs in *N. randalli* populations of Antalya Gulf. Maximum ages were III for female

and IV for male in this study. This differences in the age distribution of the populations may be due mainly to ecology of species, sampling differences, fishing activity, trophic status and ecological characteristics of the marine systems. The age composition in previous studies on species of *N. randalli* populations was as follows: Iskenderun Bay, 0-III<sup>[9]</sup> and Arabian Sea of Oman, I-II<sup>[2]</sup>. The age structure found in the population of the Antalya Gulf is different from that reported in the Iskenderun Bay and Arabian Sea of Oman.

Males attained larger length and weight than females. There are few papers from previous studies about growth parameters in *N. randalli* for comparing with results of the present study. Available literature data about the length-length relationships and morphometrics characters of *N. randalli* are very scarce<sup>[20]</sup>. The ratio of standard length:body depth, standard length:head length and head length:body depth are similar to results of Lelli *et al.*, 2008<sup>[20]</sup>.

The slope (b) value of the length weight regression of *N. randalli* populations was 3.0426. The values of b were within the limits of 2.5-3.5 commonly reported for teleosts by Froese (2006)<sup>[10]</sup>. The slope b value of the LWR equations showed a great variation from one population to another within the same

species and species of the same genus (Table 3). The  $b$  value of *Nemipterus* species in the world has been shown to range from 2.63 to 3.28. Geographic location and associated environmental conditions, such as seasonality (date and time of capture),

stomach fullness, disease and parasite loads, can also affect the value of  $b$  (Le Cren, 1951; Bagenal and Tesch, 1978)<sup>[19,3]</sup>. This study also provides a new maximum length for *N. randalli* Turkish Coast of Mediterranean.

**Table 3:** Records of *Nemipterus* specimens from the world and showing some growth values

Species	Locality	L <sub>MIN</sub>	L <sub>MAX</sub>	a	b	R <sup>2</sup>	References
<i>N. bathybius</i>	Beibu Gulf. Northern South China Sea	4.2	24.0	0.0353	2.88	0.999	Wang <i>et al.</i> 2011
<i>N. bipunctatus</i>	Chennai Fisheries Harbour			0.0000096	3.051	0.954	Kizhakudan and Rajapackiam. 2011
<i>N. furcosus</i>	Beibu Gulf. Northern South China Sea	6.20	22.2	0.0237	3.02	0.999	Wang <i>et al.</i> 2011
<i>N. hexodon</i>	Beibu Gulf. Northern South China Sea	5.20	13.3	0.0354	2.88	0.961	Wang <i>et al.</i> 2011
<i>N. hexodon</i>	Thailand Gulf	13.10	21.7	0.00576	3.277	0.99	Yanagawa. 1988
<i>N. japonicus</i>	Karnataka State	9.50	30.8	0.039	2.664	0.99	Zacharia. 1998
<i>N. japonicus</i>	Arabian Sea			0.0407	2.765	0.96	McIlwain <i>et al.</i> 2006
<i>N. japonicus</i>	Ratnagiri coast				2.629	0.9870	Kumar <i>et al.</i> 2011
<i>N. japonicus</i>	Persian Gulf. Iran	7.30	24.5	0.0517	2.664	0.928	Raeisi <i>et al.</i> 2012
<i>N. japonicus</i>	Chennai Fisheries Harbour			0.0000308	2.843	0.934	Kizhakudan and Rajapackiam. 2011
<i>N. japonicus</i>	Beibu Gulf. Northern South China Sea	4.0	20.50	0.0329	2.94	0.999	Wang <i>et al.</i> 2011
<i>N. mesoprion</i>	Chennai Fisheries Harbour			0.0000125	3.014	0.933	Kizhakudan and Rajapackiam. 2011
<i>N. metopias</i>	Beibu Gulf. Northern South China Sea	5.60	22.5	0.0384	2.84	0.999	Wang <i>et al.</i> 2011
<i>N. nematophorus</i>	Chennai Fisheries Harbour			0.000032	2.814	0.960	Kizhakudan and Rajapackiam. 2011
<i>N. nemurus</i>	Beibu Gulf. Northern South China Sea	6.50	19.8	0.0263	2.97	0.988	Wang <i>et al.</i> 2011
<i>N. peronii</i>	Coral Reef-Lagoons of new Caledonia	16	24.50	0.0157	3.029		Letourneur <i>et al.</i> 1998
<i>N. peronii</i>	Chennai Fisheries Harbour			0.0000279	2.835	0.955	Kizhakudan and Rajapackiam. 2011
<i>N. randalli</i>	Israel continental shelf	4	23.5	0.0101	3.08	0.97	Edelist. 2014
<i>N. randalli</i>	Iskenderun Bay	4.6	15.3	0.013	2.687	0.979	Erguden <i>et al.</i> 2009
<i>N. randalli</i>	Iskenderun Bay	4.8	21.5	0.0011	3.061	0.982	Erguden <i>et al.</i> 2010
<i>N. randalli</i>	Antalya Gulf	9.50	22	0.0120	2.975	0.937	Ozvarol. 2014
<i>N. virgatus</i>	Beibu Gulf. Northern South China Sea	4.30	31.6	0.0263	2.97	0.999	Wang <i>et al.</i> 2011
<i>N. zysron</i>	Chennai Fisheries Harbour			0.0000132	2.984	0.964	Kizhakudan and Rajapackiam. 2011
<i>N. zysron</i>	Coral Reef-Lagoons of new Caledonia	11.50	27	0.0103	3.167		Letourneur <i>et al.</i> 1998
<i>N. randalli</i>	Antalya Gulf	6	24	0.0105	3.0426	0.983	This study

The differences obtained in the growth parameters  $k$  and  $L_{\infty}$  between the populations of *N. randalli* could be explained on the basis of different oceanographic conditions, sampling differences and age readings. The phi-prime test estimations provide an indicator of the reliability of estimates since it is suggested that values are similar for the same species and genera. The phi-prime test estimation was calculated as 2.38 in the *N. randalli* population and the phi-prime test estimation values for the *Nemipterus* genus ranged from 2.42 to 2.97. These growth performance variations could be caused by different results obtained in age readings by the different researchers. However, it is possible that variations in population parameters of the genus *Nemipterus* represent epigenetic responses to the different environmental conditions, such as temperature, pH, dissolved oxygen levels, and geographic location. Nutrient levels also probably vary between study areas.

The rate of fish invasions into the Mediterranean Sea has increased in recent decades. They have collectively significant ecological and economic impacts on the Eastern Mediterranean fish populations<sup>[7]</sup>. Threadfin bream, *N. randalli*, become an important fisheries resources for small trawlers, especially coast

of Antalya Gulf. In conclusion, this study provides some basic information about one of the Lessepsian migrant species and is thought to be useful for future studies. The role of *N. randalli* within the coastal ecosystem and its effect on local populations need to be the subject of future research.

## 5. Acknowledgements

We would like to thank Sevilyay Karatas and Fatih Fikret Oztekin for their kind help during the field survey and laboratory studies.

## 6. References

1. Ali M, Saad A. First Records of Randall's Threadfin Bream *Nemipterus randalli* (Osteichthyes: Nemipteridae) off The Syrian Coast (Eastern Mediterranean). *Annales Series Historia Naturalis* 2013; 23(2):119-124.
2. Al-Kiyumi F, Mehanna S, Al-Bulush N. Growth, mortality and yield per recruit of the Randall's threadfin bream *Nemipterus randalli* (Russell, 1986) from the Arabian Sea off Oman. *Thalassas* 2014; 30(1):67-73.
3. Bagenal TB, Tesch F. Age and Growth. In: *Methods for assessment of fish production in fresh waters*. T. B. Bagenal

- (Ed.) Blackwell Scientific Publications, London. IBH Handbook 1978; 3:101-136.
4. Bohlen J, Freyhof J, Nolte A. Sex ratio and body size in *Cobitis elongatoides* and *Sabanejewia balcanica* (Cypriniformes, Cobitidae) from a thermal spring. *Folia Zoologica* 2008; 57:191-197.
  5. Bilecenoglu M, Russell BC. Record of *Nemipterus randalli* Russell, 1986 (Nemipteridae) from Iskenderun Bay, Turkey. *Cybiurn* 2008; 32:279-280.
  6. Edelist D. New length–weight relationships and Lmax values for fishes from the Southeastern Mediterranean Sea. *Journal of Applied Ichthyology* 2014; 30:521-526.
  7. ElHaweet AEA. Biological studies of the invasive species *Nemipterus japonicus* (Bloch, 1791) as a Red Sea immigrant into the Mediterranean. *Egyptian Journal of Aquatic Research* 2013; 39:267-274.
  8. Erguden D, Turan C, Gurlek M. Short Communication Weight–length relationships for 20 Lessepsian fish species caught by bottom trawl on the coast of Iskenderun Bay (NE Mediterranean Sea, Turkey). *Journal of Applied Ichthyology* 2009; 25:133-135.
  9. Erguden D, Turan C, Gurlek M, Yaglioglu D, Gungor M. Age and growth of the Randall's threadfin bream *Nemipterus randalli* (Russell, 1986), a recent Lessepsian migrant in Iskenderun Bay, northeastern Mediterranean. *Journal of Applied Ichthyology* 2010; 26:441-444.
  10. Froese R. Cube law, condition factor and weight–length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology* 2006; 22:241-253.
  11. Froese R, Tsikliras AC, Stergiou KI, Editorial note on Weight-Length relations of fishes. *Acta Ichthyologica et Piscatoria* 2011; 41(4):261-263.
  12. Froese R, Pauly D. FishBase. 2015. <http://www.fishbase.org> (accessed 20 January 2015)
  13. Gokoglu M, Guven O, Balci BA, Colak H, Golani D. First records of *Nemichthys scolopaceus* and *Nemipterus randalli* and second record of *Apterichthys caecus* from Antalya Bay, Southern Turkey *Marine Biodiversity Records* 2009; 6234:1-3.
  14. Golani D, Sonin A. The Japanese threadfin bream *Nemipterus japonicus*, a new Indo-Pacific fish in the Mediterranean Sea. *Journal of Fish Biology* 2006; 68:940-943.
  15. Gulsahin A, Kara A. Record of *Nemipterus randalli* Russell, 1986 from the southern Aegean Sea (Gökova Bay, Turkey). *Journal of Applied Ichthyology* 2013; 29:933-934.
  16. Gurlek M, Erguden S, Yaglioglu D, Turan F, Demirhan S, Gurlek M *et al.* Feeding Habits of Indo-Pacific Species *Nemipterus randalli* Russel, 1986 (Nemipteridae) in Iskenderun Bay, Eastern Mediterranean Sea. *Rapp Comm int Mer Médit* 2010; 39:539.
  17. Kizhakudan SJ, Rajapackiam S. Length-weight relationship in six species of threadfin breams occurring in the trawl landings at Chennai. *Journal of the Marine Biological Association of India* 2011; 53(2):268-271.
  18. Kumar PS, Mohite SA, Naik SD, Mohite AS. Length weight relationship in *Nemipterus japonicus* of Ratnagiri coast along Maharashtra. *Indian Journal of Applied and Pure Biology* 2011; 26(1):79-84.
  19. LeCren ED. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology* 1951; 20:201-219.
  20. Lelli S, Colloca F, Carpentieri P, Russell BC. The threadfin bream *Nemipterus randalli* (Perciformes: Nemipteridae) in the eastern Mediterranean Sea. *Journal of Fish Biology* 2008; 73:740-745.
  21. Letourneur Y, Kulbicki M, Labrosse P. Length-weight Relationship of Fishes from Coral Reefs and Lagoons of New Caledonia - An Update. *Naga, the ICLARM Quarterly*, 1998, 39-46.
  22. McIlwain J, Hermosa GV, Claereboudt M, Al-Oufi HS, Al-Awi M. Spawning and Reproductive patterns of six exploited finfish species from the Arabian Sea, Sultanate of Oman. *Journal of Applied Ichthyology* 2006; 22:167-176.
  23. Ozvarol Y. Length–weight relationships of 14 fish species from the Gulf of Antalya (northeastern Mediterranean Sea, Turkey). *Turkish Journal of Zoology* 2014; 38:342-346.
  24. Pauly D. Fish population dynamics in tropical waters: a manual for use with programmable calculators. *ICLARM Studies and Reviews* 8, International Center for Living Aquatic Resources Management, Philippines, Manila, 1984, 325.
  25. Pauly D, Munro JL. Once more on the comparison of growth in fish and invertebrates. *Fishbyte* 1984; 2:21.
  26. Raeisi H, Paighambari SY, Davoodi R, Bibak M, Hoseini, SA, Shabni MJ. Length-weights relationships and relative weights of some demersal fish species from the Persian Gulf, Iran. *African Journal of Agricultural Research* 2012; 7(5):741-746.
  27. Russell BC. Nemipterid fishes of the world (threadfin breams, whiptail breams, monocle breams, dwarf monocle breams, and coral breams). Family Nemipteridae. An annotated and illustrated catalogue of nemipterid species known to date. *FAO Fisheries Synopsis* no. FAO, Rome. 1990; 125:12.
  28. Russell BC. A review of the threadfin breams of the genus *Nemipterus* (Nemipteridae) from Japan and Taiwan, with description of a new species. *Japanese Journal of Ichthyology* 1993; 39:295-310.
  29. Wang XH, Qiu YS, Zhu GP, Du FY, Sun DR, Huang SL. Technical contribution Length-weight relationships of 69 fish species in the Beibu Gulf, northern South China Sea. *Journal of Applied Ichthyology* 2011; 27:959-961.
  30. Yanagawa H. Length-Weight Relationship of Gulf of Thailand Fishes. *Naga, the ICLARM Quarterly* 1988; 17(4):48-52.
  31. Zacharia PU. Dynamics of the thread-fin bream, *Nemipterus japonicus* exploited off Karnataka. *The Indian Journal of Fisheries* 1998; 45(3):265-270.