



International Journal of Fisheries and Aquatic Studies

ISSN: 2347-5129

IJFAS 2014; 2(1): 244-247

© 2013 IJFAS

www.fisheriesjournal.com

Received: 10-07-2014

Accepted: 05-08-2014

Shahlina Haque

Department of Life Sciences,
Dibrugarh University,
Dibrugarh, Assam, India.

S.P. Biswas

Department of Life Sciences,
Dibrugarh University,
Dibrugarh, Assam, India.

Length-Weight relationship and condition factor of *Botia dario* (Hamilton-Buchanan) from Sivasagar district

Shahlina Haque and S.P. Biswas

Abstract

The length-weight relationship and condition factor of *Botia dario* were carried out from wetland of Sivasagar District of Assam between March'2012 and February'2014. This paper throws light on the changes in the growth coefficient (b) seasonally and condition factor. Length-weight relationship and condition factor for a sample size of 300 specimens were calculated on seasonal basis. A wide fluctuation was observed in growth coefficient. The 'b' value ranged from 3.06 in male and 3.128 in female length-wise. Seasonally, 'b' value for male is 2.02 to 3.45 and for female is 2.4 to 3.17.

Keywords: Length-weight relationship, *Botia dario*, Condition factor.

1. Introduction

Length-weight relationship is of great considerable importance in fishery research especially for study of fish population dynamics and stock condition [7, 14, 17]. According to [21], this relationship is useful to estimate the relative condition factor to assess the general well being of the fish or type of somatic growth whether isometric or allometric. These studies are mainly directed towards two objectives; first to establish a mathematical relationship between two variables; length and weight and secondly to know the variations from expected weight for various length groups [25]. Very little works were done by [9, 4, 8, 23].

Condition factor reflects the fluctuations by interaction among feeding conditions, parasitic infection and physiological factors and recent physical and biological circumstances [21]. The study of condition assumes that heavier organisms of a given length are in better physical state, therefore, condition indices are used in fishery sciences as indicators of the length-weight-relationship of a population. Condition factor also helps to reflect the feeding conditions of the species. The condition factor, in fisheries science, is used in order to compare the "condition", "fitness" or well being of fish. It is based on the hypothesis that heavier fish of a particular length are in a better physiological condition [7].

Botia dario (Hamilton) is a small sized (maximum recorded length 10cm) loach found in streams with sandy bottom as well as in wetlands. It is one of the highly priced aquarium fish of the Brahmaputra basin. It naturally breeds in monsoon season [10]. The present communication deals with the length-weight relationship and condition factor of *Botia dario* from Sivasagar, Assam.

2. Material and Methods

Samples were collected from the selected wetlands of Sivasagar district between March'2012 and February'2014. The collected specimens were preserved in 5% formalin. The specimens' length were measured to the nearest cm and weighted to the nearest gm. The length-weight relationship was based on 250 specimens collected during study period.

Length-weight relationship was estimated by using the equation $W=aL^b$ (LeCren, 1951), where W=weight, L=length. 'a' and 'b' are initial growth and growth coefficient respectively. The values of constant 'a' and 'b' were estimated from coefficient of regression equation. The correlation coefficient (r) was estimated to determine the degree of linear

Correspondence:

Shahlina Haque

Department of Life Sciences,
Dibrugarh University,
Dibrugarh, Assam, India.

relationship between the length and weight samples. Condition factor or ponderal index was calculated by using formula $K=100W/L^3$ [6] to estimate the general wellbeing of the fish.

3. Result and Discussion

The average total length of male is 6.5 cm. and that of female is 8.7 cm. The average total weight for male and female are 3.8 and 8.9 respectively. The length-weight relationship

obtained for the males and females of *Botia dario* in different seasons have been shown in Table 1. A linear relationship between the length and weight as well as condition factor was established. The regression equation represented for male and female are as follows:

Males: $\text{LogW}=\text{Log } 0.663 + 3.06 \text{ LogL}$

Females: $\text{LogW}=\text{Log } -2.076 + 3.28\text{LogL}$

Table 1: Seasonal variation of length-weight relationship and condition factor of *Botia dario*

Season	Sex	Length Mean + SD	Weight Mean + SD	b	r	LogW=log a+b logL	k
Pre-Monsoon	M	8.7 ± 1.84	8.4 ± 4.4	3.45	0.99	log -2.4 + 3.45logL	1.11
	F	8.3 ± 1.3	7.2 ± 2.8	3.17	0.96	log -2.09 + 3.17logL	1.18
Monsoon	M	6.16 ± 2.05	2.9 ± 2	3.09	0.88	log -2.03 + 3.09logL	1.05
	F	11.45 ± 1.83	19.9 ± 8.13	3.01	0.98	log -1.92 + 3.01logL	1.25
Post-Monsoon	M	5.5 ± 1.15	2.1 ± 1.6	2.5	0.87	log -1.6 + 2.5logL	1.03
	F	7.34 ± 0.85	4.3 ± 1.5	2.5	0.86	log -1.64 + 2.5logL	1.1
Winter	M	5.64 ± 0.95	2.1 ± 1.16	2.02	0.83	log -1.2 + 2.02logL	1.085
	F	7.7 ± 1.00	4.3 ± 1.5	2.4	0.9	log -1.5 + 2.4logL	0.96




Fig 1: Length-weight relationship in *B. dario* (female)




Fig 2: Length-weight relationship in *B. dario* (male)

The length-weight relationship was calculated for different seasons to see seasonal fluctuation. Four seasons viz., winter (Dec-Feb), pre monsoon (Mar-May), monsoon (Jun-Aug) and post monsoon (Sept-Nov) have been taken into consideration. In *Botia dario* the growth coefficient (coefficient for regression) for different seasons was found

within the range of 2.02 (winter) to 3.45 (pre-monsoon) for male and from 2.4 (winter) to 3.17 (pre-monsoon) for female. The coefficient of correlation for different seasons was found to range from 0.83 to 0.99 for male and from 0.86 to 0.98 for female.

According to [21], the variation in ‘b’ value is due to

environmental factors, seasons, food availability, sex, life stage and other physiological factors. It is evident from the results that the value of regression coefficient 'b' of *Botia dario* tends to be higher during pre-monsoon and lower during winter. The 'b' value may change seasonally and even daily and also between habitats [7, 16]. A characteristic length-weight relationship for fishes is that the value of exponent (b) is 3 when growth is said to be isometric. If 'b' value is different from 3, growth is said to be allometric. The variation in fish sizes indicate that the fish population ranged from immature specimens to fully matured ones. This also suggest differences in their growth [13]. The b value 3 is not confined for all fishes because growth causes for the change of their shapes [2]. In the present study higher regression coefficient ($b > 3$) was observed for both the sexes indicating a positive allometric growth during pre-monsoon and monsoon. However, lower value of regression coefficient ($b < 3$) were recorded in post-monsoon and winter season was an indication of negative allometric growth. The fish did not follow the cube law ($b = 3$) strictly. The regression coefficient (b) of female was higher than that of the males. Higher regression value in female was also obtained by [5] in *Labeo bata* from Bangladesh; [12] reported the same in *Lepidocephalichthys guntea* from Morang District, Assam; [22] in case of *Puntius sophore* from flood plain wetland in West Bengal; [18] in *Rasbora daniconius* from Karnataka; [24] in *Puntius filamentous* from Chalakudy River, Kerala; [8] in *Amblypharyngodon mola* in Garjan beel of Assam; [15] in *Amblypharyngodon mola* from Assam.

The correlation coefficient (r) values suggested that the length and weight of female (0.93) are slightly better correlated than the male (0.89). As a whole, the maximum and minimum value of 'r' in female of *B. dario* was recorded in monsoon and post-monsoon, while in male the maximum and minimum values were recorded in pre-monsoon and winter respectively. In fish, the weight is considered to be a function of length [27]. Further, [28] viewed that exact length-weight relationship differs among fish species according to their inherited body shape and within a species according to the condition of individuals, availability of food and prior to sampling growth within the weeks.

The condition factor of *B. dario* was found to range from 1.03 to 1.11 in males and 0.96 to 1.25 in females seasonally. The highest condition value was observed during monsoon for female and pre-monsoon for males. It has been observed that females were found to be slightly in better condition than the male. Similar observation have also been reported by many workers [1, 19, 11, 23]. Biotic and abiotic environmental factors highly influenced the condition factor, and the index can be used to assess the status of the aquatic ecosystem in which fish live [3]. Through the variation of condition factor, it reflects information on physiological state of the fish on its welfare [20]. K also gives information when comparing two populations living in certain feeding, density, climate and other conditions; when determining the period of gonad maturation; and when following up the degree of feeding activity of a species to verify whether it is making good use of its feeding source [26].

4. Acknowledgement

The authors are grateful to the Department of Life Sciences, Dibrugarh University for providing necessary facilities to carry out the work. Also thankful to Zoology Department of Sibsagar College for permitting me to use the laboratory.

5. References

1. Alam MM, Rahman T, Parween S. Morphometric characters and condition factors of five freshwater fishes from Pagla river of Bangladesh. *Int J Aqu Bio* 2014; 2(1):14-19
2. Ali SS. *Fresh Water Fishery Biology*. Naseem Book Depot, Hyderabad, 1999, 330.
3. Anene A. Condition factors of four cichlid species of a man-made lake in Imo state, Southeast, Nigeria. *Turk J Fish Aquat Sci* 2005; 5:43-47.
4. Anna MTV, Jacob E, Bhaskar RK. Length-weight relationship of sixteen species of indigenous ornamental fishes of the Western Ghats of India. *Indian J Fish* 2008; 55(4):337-339.
5. Azadi MA, Naser A. Length-weight relationship and relative condition factor of a carp, *Labeo bata* (Ham.) from Kaptai Reservoir, Bangladesh. *Chittagong Univ Stud Part II: Sci* 1996; 20(2):19-25.
6. Bagenal TB. Aspects of fish fecundity. In: S.D. Gerking (Ed) *Ecology of Freshwater fish Production*. Blackwell Scientific Publications Oxford: 1978; 75-101.
7. Bagenal TB, Tesch FW. Age and Growth. In: Bagenal, T. (Ed.), *Methods of Assessment of fish Production in fresh waters*. Oxford Blackwell Scientific Publication. 1978; 101-136.
8. Baishya A, Dutta A, Bordoloi S. Morphometry and length-weight relationship of *Amblypharyngodon mola* (Hamilton-Buchanan, 1822). In *Indian J Fish* 2010; 57(1):87-91.
9. Chandrika B, Balasubramonian NK. Length-weight relationship of *Xenentodon cancila* (Ham.) (Teleostei: Belontiidae). *Proc India Acad Sci* 1986; 95(2):187-190
10. Das JN, Biswas SP. *A Handbook of Ornamental Fishes of the Brahmaputra Basin*. EBH Publishers (India), Guwahati, 2008; 109.
11. Dasgupta M. Length-weight relationship and condition factor (K value) for *Tor putitora* (Ham.) from Garo Hills, Meghalaya. *Indian J Fish* 1991; 38(1):35-38
12. Dhakal A, Subba BR. Length-weight relationship of *Lepidocephalichthys guntea* of Pathri Khola, Morang District. *Our Nature* 2003; 1:53-57.
13. Forta LO, Costa PAS, Braga AC. Length-weight relationship of marine fishes from the central Brazilian coast. *NAGA, ICLARMQ* 2004; 27(1&2):20-26.
14. Garcia CB, Buarte JO, Sandoval N, Von Schiller D, Mello, Najavas P. Length weight relationships of Demersal Fishes from the Gulf of Salamanca, Colombia. *Fishbyte* 1998; 21:30-32.
15. Gogoi R, Goswami UC. Length-weight relationship and sex ratio of freshwater fish *Amblypharyngodon mola* (HAM-BUCH) from Assam. *Int J Fish Aqu Stu* 2014; 1(4):68-71.
16. Gonclaves JMS, Bente L, Lino PG, Ribeiro J, Canario AVM, Erzini K *et al*. Weight-length relationships for selected fish species of the small-scale demersal fisheries of the south and south-west coast of Portugal. *Fish Res* 1997; 30:253-256.
17. Haimovici M, Velasco G. Length-weight relationship of marine fishes from southern Brazil. *The ICLARM Quarterly* 2000; 23(1):14-16.
18. Kumar HK, Kiran BR, Purushotham R, Puttaiah ET, Manjappa S. Length-weight relationship of cyprinid fish *Rasbora daniconius* (Hamilton-Buchanan) from Sharavathi reservoir, Karnataka. *Zoos' Print Journal*

- 2006; 21(1):2140-2141.
19. Jhingran VG. General Length-weight relationship of three major carps of India. Proc Nat Inst Sci India 1952; 18:449-460.
 20. Kumolu-Johnson CA, Ndimele PE. Length-weight relationships and Condition Factors of Twenty-One Fish species of Ologe Lagoon, Lagos, Nigeria. Asian J Agricultural Sciences 2010; 2(4):174-179.
 21. LeCren ED. The length-weight relationship seasonal cycle in gonadal weight and condition factor of perch (*Perca fluviatilis*). J Anim Ecol 1951; 20:201-219.
 22. Mitra K, Suresh VR, Vinci GK, Naskar B. Length-weight relation, reproductive characters and condition of *Puntius sophore* (Hamilton) from a flood plain wetland in West Bengal. Inland Fish Soc India. 2005; 37(1):16-22.
 23. Paswan G, Abujam SKS, Dey M, Biswas SP. Length-weight relationship of two species of *Trichogaster* (Colisa) from Brahmaputra Basin of Assam, J Bio Innov 2012; 1(1):6-13
 24. Prasad G, Ali APH. Length-weight relationship of a cyprinid fish *Puntius filamentosus* from Chalakudy River, Kerala. Zoos Print 2007; 22(3):2637-2638.
 25. Srivastava R, Singh HR. Length-weight Relationship of *Puntius sophore* (Ham.) from Allahabad region. Proc Nat Acad Sci India 2003; 73B(1):53-58.
 26. Weatherley AH. Growth and Ecology of fish populations. Academic Press, London. 1972; 293.
 27. Weatherley AH, Gill HS. The biology of fish growth. Academic Press, London, 1987, 443.
 28. Yousuf F, Khurshid S. Length-weight relationship and relative conditions factor for the halfbeak *Hemiramphus far* (Forsskal, 1775) from the Karachi coast. Univ J Zool Rajshahi Univ 2008; 27:103-104.