



International Journal of Fisheries and Aquatic Studies

E-ISSN: 2347-5129
P-ISSN: 2394-0506
(ICV-Poland) Impact Value: 5.62
(GIF) Impact Factor: 0.549
IJFAS 2017; 5(1): 403-406
© 2017 IJFAS
www.fisheriesjournal.com
Received: 09-11-2016
Accepted: 10-12-2016

Chryssa Anastasiadou

1) Centre for Ecological and
Evolutionary Synthesis,
Department of Biosciences,
University of Oslo, Blindern,
Oslo, Norway

2) Fisheries Research Institute,
Nea Peramos, Kavala,
Macedonia, Greece

Moslem Daliri

Fisheries Research Institute, Nea
Peramos, Kavala, Macedonia,
Greece

Anthi Oikonomou

Fisheries Department, Faculty of
Marine and Atmospheric
Sciences, Hormozgan University,
Bandar Abbas, Iran

Leif Asbjørn Völlestad

Laboratory of Zoology,
Department of Biological
Applications and Technology,
University of Ioannina,
University Campus, Ioannina,
Epirus, Greece

Correspondence

Chryssa Anastasiadou

1) Centre for Ecological and
Evolutionary Synthesis,
Department of Biosciences,
University of Oslo, Blindern,
Oslo, Norway

2) Fisheries Research Institute,
Nea Peramos, Kavala,
Macedonia, Greece

Length weight relationships, relative weight and relative condition factor of three freshwater shrimps from Greece

Chryssa Anastasiadou, Moslem Daliri, Anthi Oikonomou and Leif
Asbjørn Völlestad

Abstract

Length-weight relationships (LWR), relative weight and relative condition factor were studied for three freshwater carideans (*Atyaephyra acheronensis*, *Atyaephyra thymisensis* and *Palaemon antennarius*) from lotic and lentic habitats of northwestern Greece. The two atyids are Adriatic-Ionian endemics while the palaemonid is distributed in the Apennine Peninsula, Sardinia, Sicily and Adriatic-Ionian eco regions. Data were collected from 760 individuals and LWR parameters were estimated by the least-squares method. Kolmogorov-Smirnov and Levene tests were used to analyse data normality and homogeneity of variances. The b values ranged from 2.908 to 4.134. More specifically, in *A. acheronensis*, R^2 ranged from 0.93 to 0.96, (b values = 2.983 - 3.673), in *A. thymisensis*, R^2 ranged from 0.61 to 0.91, (b values = 3.489 - 3.714), and in *P. antennarius* populations, R^2 ranged from 0.47 to 0.95 (b values = 2.908 - 4.135). The relative weight showed no significant differences between sexes and species for both *A. thymisensis* and *A. acheronensis* while for *P. antennarius* ranged from 53.07% to 93.56% for males and from 66.78% to 85.82 for females among the studied sites. The relative condition factor in *A. thymisensis* males found 0.9 ± 0.02 and in females 1 ± 0.02 ($P < 0.05$) while in *A. acheronensis*, showed no differences between sexes. The same parameter ranged from 0.84 ± 0.02 to 1.06 ± 0.02 and from 0.79 ± 0.02 to 1.06 ± 0.02 for *P. antennarius* males and females, respectively. The present study contributes to the first recording on the LWR for the studied species, providing basic and valuable information for shrimp fisheries management.

Keywords: Caridea, decapoda, atyidae, palaemonidae, morpho-gravimetric data, allometric growth

1. Introduction

In order to properly evaluate a fishery information on the growth processes is important to record the major morphometric and gravimetric parameters of the individuals. The growth processes of the different organisms can be assessed in various ways. One alternative way is to describe how the body mass changes with various length measures. Such length-weight relationships (LWRs) are commonly used as predictive scientific tools in fisheries (Petrakis and Stergiou, 1995; Vaslet *et al.*, 2008) [20, 24]. Knowledge on LWRs may be important for biomass estimation, evaluation of the stock condition, and when forecasting stock development. Such information is needed in order to properly manage marine and freshwater ecosystems (Ecoutin *et al.*, 2005, Daliri *et al.*, 2012a, b) [13, 9, 10]. Despite the importance of having access to LWRs, to date, little such information is available for shrimps. This is unfortunate as there is large global interest for harvested (Chu *et al.*, 1995; Guéguen, 1997; Guijarro *et al.*, 2012) [8, 16, 17] and cultured marine shrimps (Araneda *et al.*, 2008) [2] and also to a lesser degree for estuarine/freshwater shrimps (Enin, 1994; Brouwer *et al.*, 2007; Deekae and Abowei, 2010) [14, 4, 11]. More information is thus needed.

The recent revisions of two freshwater natant decapod genera *Atyaephyra* de Brito Capello, 1867 and *Palaemon* Weber, 1795 (Christodoulou *et al.*, 2012; Tzomos and Koukouras, 2015; Carvalho *et al.*, 2014; Christodoulou *et al.*, 2016) [6, 23, 5, 7] along with the descriptions of six new species revealed a high Balkan endemism in the Mediterranean region. Seven species have been described within the genus *Atyaephyra*, four of which are Balkan endemics (*A. acheronensis*, *A. stankoi*, *A. strymonensis* and *A. thymisensis*) (Christodoulou *et al.*, 2012) [6], while nine *Palaemon* species have been reported from the area with *P. antennarius* being

distributed only in the Apennine Peninsula, Sardinia, Sicily and Adriatic-Ionian ecoregions (Tzomos and Koukouras, 2015) [23]. Although there is published information on the bionomy of various species of the two genera (Christodoulou *et al.*, 2016 and references herein) [7], LWRs have not been published. Further, for crustaceans such as shrimps it is not evident which length metrics are best suited for describing LWRs. Based on the above, the present study aims at: 1) reporting the LWRs, the relative weight, and the relative condition factor of three species (*Atyaephyra acheronensis*, *Atyaephyra thymisensis* and *Palaemon antennarius*), and 2) evaluating three different body lengths in order to optimize the LWR analyses.

2. Materials and Methods

Shrimp individuals were collected during April 2004 from four sites of NW Greece [S1: Lake Pamvotis (39.3920° N, 20.5214° E), S2: Louros River (39.0314° N, 20.4626° E), S3: Acheron River (39.1396° N, 20.2914° E), S4: Thyamis River (39.3228° N, 20.0976° E)] by means of a handnet with a mesh size of 2mm and preserved *in situ* with 4% formalin solution. Species and sex were determined via stereo-microscopic inspection (LEICA MZ12^s). Gravimetric data were collected by means of a digital caliper (0.01 mm; Vogel DigiPlus) and a digital scale (0.01 g; Ohaus-Adventurer). Three body length measurements were taken on each individual: 1) CL: distance from the orbit to the posterior point of carapace, 2) TL1: distance from the orbit to the posterior point of sixth somite and 3) TL2: distance from the tip of rostrum to the posterior end of the uropodal exopod. LWRs were estimated by the equation $W = aL^b$, where W is the total weight in g, L is the measured length (CL, TL1 and TL2) in mm, a is the intercept, and b is the exponent. Parameters a and b were estimated by the least-squares method based on the logarithmic equation: $\log W = \log a + b \log L$. For each specimen, Relative Weight (RW) and Relative Condition Factor (RK) were calculated using the following equations respectively:

a) $RW = 100 (W/W_s)$, where W is the weight of a specimen and W_s is the standard weight representing the 75th percentile of observed weights at that length (Froese, 2006) [15]

b) $RK = W/aL^b$, where W is the body weight, L is CL, and a and b are the parameters of the LWR (Le Cren, 1951) [18].

Kolmogorov-Smirnov and Levene tests were used to analyse for normality of the data and homogeneity of variances. Two independent samples t-tests (5% level) were used for the comparison of the condition of males and females individuals in each ecosystem. Oviparous females were excluded from analysis, so that the maturation state of the egg masses could not influence the comparisons. All the analyses were conducted using SPSS 11.

3. Results and Discussion

Overall, 760 individuals were collected, measured and analyzed. The results of the LWR regressions, and the relative weight and relative condition factor estimates are presented in Table 1. Among the studied species, CL ranged from 3.6 to 11.8mm, TL1 from 11.7 to 32.9mm, TL2 from 17.2 to 46.6mm and TW from 0.028 to 0.981g (Tab. 1). *A. acheronensis* and *A. thymisensis* showed isometric ($b \approx 3$) to positive allometric ($b > 3$) growth ($b = 2.983$ and $b = 3.489$, respectively), a fact that has been previously been shown for *Macrobrachium macrobrachion*, especially from estuarine ecosystems (Enin, 1994) [14]. High b values are usually correlated with eutrophic conditions and abundant food (Froese, 2006, Liasko *et al.*, 2011) [15, 19].

Among the studied ecosystems (denoted S1-S4), *P. antennarius* showed isometric [b (S4) = 2.908] to positive allometric growth patterns [b (S1) = 4.135, b (S2) = 3.477 and b (S3) = 3.294], with R^2 ranging from 0.79 to 0.95. Only the *P. antennarius* population from Acheron River (S3) demonstrated low R^2 values (Tab. 1). Similar b values have been reported for *Macrobrachium acanthurus*, *M. olfersii*, *Palaemon northropi* (Anger and Moreira, 1998) [1], *Palaemon serratus* (Reeve, 1969) [21], *Palaemonetes argentinus* (Donatti, 1986) [12], *M. vollenhovenii* (Willführ-Nast *et al.*, 1993) [26], *M. rosenbergii* (Wickins, 1972) [25] and *M. lamarrei* (Shafi *et al.*, 1977) [22]. For the three selected lengths, the best fit for LWR was given by TL2 measurement, as has also been proposed by Anger and Moreira (1998) [1].

The relative weights (RW), which is the most valuable factor for comparative studies (Blackwell *et al.*, 2000; Daliri *et al.*, 2012a, b) [3, 9, 10], differences between the sexes in *P. antennarius* populations (Kruskal-Wallis test, $\chi^2 = 14.035$ and $P < 0.001$) (Tab. 1). The Mann-Witney U test ($\alpha = 0.5$) showed that the conditions are favorable for females in Lake Pamvotis (S1) and for males in Louros River (S2). Moreover, the data of the present study reveal that *A. acheronensis* had better performance (RW=89%) than *A. thymisensis* (RW = 68%). A general linear model (GLM) analysis which was used to analyze the effect of the three factors (sex, type of species and ecosystem with interactions) on RW, showed a statistically significant interaction between the type of species and sex ($P < 0.001$). Finally, RK showed almost the same values for *Atyaephyra* species while the lowest values were observed for *P. antennarius* from the lentic habitat (S1) (Tab. 1).

This study constitutes the first report on the LWRs for 3 freshwater shrimps (*Atyaephyra acheronensis*, *Atyaephyra thymisensis* and *Palaemon antennarius*) two of which (atyids) are Balkan endemics. Moreover, the relative weights and the relative condition factors of the studied species are given for the first time and comparatively for the four main ecosystems of north-western Greece.

Table 1: Calculated parameters of LWR, relative weights and relative condition factors for the studied species. NI: number of individuals, CL: carapace length, TL1: total length 1, TL2: total length 2, TW: total weight. Study sites: S1, S2, S3, S4 (are detailed in Material and Methods).

Species (Site)	NI	CL (mm)		TL1 (mm)		TL2 (mm)		TW (g)	
		Min	Max	Min	Max	Min	Max	Min	Max
<i>A. acheronensis</i> (S3)	100	3.6	7.3	11.7	21.4	17.2	29.7	0.028	0.234
<i>A. thymisensis</i> (S1)	91	4.1	7.3	13.8	21.6	20.5	31.4	0.047	0.268
<i>P. antennarius</i> (S1)	273	5.1	11.8	15.4	32.9	22.6	46.6	0.069	0.981
<i>P. antennarius</i> (S2)	100	4.6	9.4	13.0	25.4	20.0	35.8	0.076	0.427
<i>P. antennarius</i> (S3)	95	4.4	9.1	12.9	26.1	19.1	35.5	0.044	0.435
<i>P. antennarius</i> (S4)	101	4.8	8.9	16.3	23.9	22.4	34.4	0.088	0.363
Length-weight parameters									
Species (Site)	Type of Length	a		b (95% CI of b)		R ²	P		
<i>A. acheronensis</i> (S3)	CL	6x10 ⁻⁴		2.983 (2.903-3.063)		0.93	<0.001		

	TL1		8x10 ⁻⁶		3.303 (3.143-3.823)	0.94	<0.001
	TL2		8x10 ⁻⁷		3.673 (3.523-3.823)	0.96	<0.001
<i>A. thymisensis</i> (S1)	CL		3x10 ⁻⁴		3.489 (3.319-3.659)	0.91	<0.001
	TL1		5x10 ⁻⁶		3.499 (2.975-4.019)	0.66	<0.001
	TL2		7x10 ⁻⁷		3.714 (3.194-4.234)	0.69	<0.001
<i>P. antennarius</i> (S1)	CL		2x10 ⁻⁴		3.591 (3.481-3.701)	0.79	<0.001
	TL1		2x10 ⁻⁶		3.746 (3.526-3.966)	0.81	<0.001
	TL2		1x10 ⁻⁷		4.135 (3.925-4.345)	0.85	<0.001
<i>P. antennarius</i> (S2)	CL		9x10 ⁻⁴		2.758 (2.628-2.888)	0.95	<0.001
	TL1		8x10 ⁻⁶		3.381 (3.171-3.627)	0.91	<0.001
	TL2		2x10 ⁻⁶		3.477 (3.327-3.627)	0.95	<0.001
<i>P. antennarius</i> (S3)	CL		9x10 ⁻⁴		2.646 (2.056-2.846)	0.46	<0.001
	TL1		3x10 ⁻⁵		2.846 (2.116-3.576)	0.39	<0.001
	TL2		2x10 ⁻⁶		3.294 (2.574-4.014)	0.47	<0.001
<i>P. antennarius</i> (S4)	CL		2x10 ⁻³		2.393 (2.173-2.613)	0.82	<0.001
	TL1		4x10 ⁻⁵		2.836 (2.586-3.086)	0.84	<0.001
	TL2		1x10 ⁻⁵		2.908 (2.688-3.128)	0.88	<0.001
Species (Site)	Sex	Ws (g)	Mean RW (%) [SE of RW]	P	Mean RK [SE of RK]	P	
<i>A. acheronensis</i> (S3)	Males	0.0587	86.34 [4.72]	<0.25	1 [0.01]	<0.788	
	Females	0.1054	88.78 [4.12]		1 [0.01]		
	Both	0.0913	79.00 [3.67]				
<i>A. thymisensis</i> (S1)	Males	0.0710	88.92 [2.23]	<0.836	0.9 [0.02]	<0.002	
	Females	0.2085	89.45 [2.48]		1 [0.02]		
	Both	0.1748	67.91 [3.96]				
<i>P. antennarius</i> (S1)	Males	0.3994	53.07 [3.22]	<0.001	0.84 [0.02]	<0.082	
	Females	0.4417	84.11 [3.55]		0.79 [0.02]		
	Both	0.3998	74.65 [2.85]				
<i>P. antennarius</i> (S2)	Males	0.1581	93.56 [5.95]	<0.001	1.06 [0.01]	<0.927	
	Females	0.3454	66.78 [5.12]		1.06 [0.02]		
	Both	0.2154	82.65 [4.46]				
<i>P. antennarius</i> (S3)	Males	0.1199	83.03 [3.52]	<0.466	0.91 [0.04]	<0.007	
	Females	0.1753	84.95 [6.20]		1.06 [0.04]		
	Both	0.1431	86.27 [4.34]				
<i>P. antennarius</i> (S4)	Males	0.1529	88.00 [3.00]	<0.077	1.02 [0.02]	<0.076	
	Females	0.2335	85.82 [5.48]		0.96 [0.02]		
	Both	0.1900	88.26 [3.98]				

4. Acknowledgement

The present study is a part of a postdoctoral research which was implemented through the project "Scholarships SSF in Integrated Marine and Inland Waters Administration" and co-funded by 85% of resources XM EOX period 2009- 2014 and by 15% from the Public Investment program.

5. References

- Anger K, Moreira GS. Morphometric and Reproductive Traits of Tropical Caridean Shrimps. *Journal of Crustacean Biology*. 1998; 18(4):823-838.
- Araneda M, Pérez PE, Gasca-Leyva E. White shrimp *Penaeus vannamei* culture in freshwater at three densities: Condition state based on length and weight. *Aquaculture*. 2008; 283(1-4):13-18.
- Blackwell BG, Brown ML, Willis DW. Relative weight (W_r) status and current use in fisheries assessment and management. *Reviews in Fisheries Science*. 2000; 8(1):1-44.
- Brouwer M, Brown-Peterson NJ, Larkin P, Patel V, Denslow N, Manning S *et al.* Molecular and whole animal responses of grass shrimp, *Palaemonetes pugio*, exposed to chronic hypoxia. *Journal of Experimental Marine Biology and Ecology*. 2007; 341(1):16-31.
- Carvalho FL, Magalhães C, Mantelatto FL. Molecular and morphological differentiation between two Miocene-divergent lineages of Amazonian shrimps, with the description of a new species (Decapoda, Palaemonidae, *Palaemon*). *Zookeys*. 2014; 457:79-108.
- Christodoulou M, Antoniou A, Magoulas A, Koukouras A. Revision of the freshwater genus *Atyaephyra* (Crustacea, Decapoda, Atyidae) based on morphological and molecular data. *ZooKeys*. 2012; 229:53-110.
- Christodoulou M, Anastasiadou Ch, Jugovic J, Tzomos Th. Freshwater Shrimps (Atyidae, Palaemonidae, Typhlocarididae) in the Broader Mediterranean Region: Distribution, Life Strategies, Threats, Conservation Challenges and Taxonomic Issues. In: A Global Overview of the Conservation of Freshwater Decapod Crustaceans. Kawai, T., Cumberlidge, N. (Eds). Springer International Publishing AG, Switzerland, 2016, 199-236.
- Chu KH, Chen QC, Huang LM, Wong CK. Morphometric analysis of commercially important penaeid shrimps from the Zhujiang estuary, China. 1995; *Fisheries Research*, 1995; 23(1-2):83-93.
- Daliri M, Eighani M, Paighambari SY, Alizadeh E. Some morphological characteristics of five marine fish species of Hormozgan coastal waters (Northern Persian Gulf). *Caspian Journal of Applied Sciences Research*, 2012a; 1(6):20-26.
- Daliri M, Paighambari SY, Shabani MJ, Pouladi M, Davoodi R. Length-weight and length-girth relationships, relative weight and relative condition factor of four commercial fish species of northern Persian Gulf. 2012a; *Annual Review & Research in Biology*, 2012a; 2(1):15-26.
- Deekae SN, Abowei JFN. *Macrobrachium macrobrachion* (Herklots, 1851) Length-weight

- relationship and Fulton's condition factor in Luubara creek, Ogoni Land, Niger Delta, Nigeria. *International Journal of Animal and Veterinary Advances*, 2010; (4):155-162.
12. Donatti OS. Algunos aspectos bioecologicos Del camaron *Palaemonetes argentinus* (Nobili, 1901) en el embalse San Roque, Cordoba (Argentina). *Revista de Hidrobiologia Tropical*, 1986; 19(1):45-60.
 13. Ecoutin JM, Albaret JJ, Trap S. Length-weight relationships for fish populations of a relatively undisturbed tropical estuary: The Gambia. *Fisheries Research*, 2005; 72(2-3):347-351.
 14. Enin UI. Length-weight parameters and condition factor of two West African prawns. *Revista de Hidrobiologia Tropical*, 1994; 27(2):121-127.
 15. Froese R. Cube law, condition factor and Length-Weight relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 2006; 22(4):241-253.
 16. Guéguen F. Morphometric relationships of two commercial deep-sea shrimp species from French Guiana. 1997; *Comptes Rendus de l'Academie des Sciences Series III Sciences de la Vie*, 1997; 320(11):899-908.
 17. Guijarro B, Fanelli E, Moranta J, Cartes JE, Massuti E. Small-scale differences in the distribution and population dynamics of pandalid shrimps in the western Mediterranean in relation to environmental factors. *Fisheries Research*, 2012; 119-120:33-47.
 18. Le Cren ED. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology*. 1951; 20(2):201-219.
 19. Liasko R, Koulis A, Pogrebniak A, Papiggioti O, Taranenko L, Leonardos I. Influence of environmental parameters on growth pattern and population structure of *Carassius auratus gibelio* in Eastern Ukraine. *Hidrobiologia*, 2011; 658(1):317-328.
 20. Petrakis G, Stergiou KI. Weight-length relationships for 33 fish species in Greek waters. *Fisheries Research*, 1995; 21(3-4):465-469.
 21. Reeve MR. *The laboratory culture of the prawn Palaemon serratus*. Fishery investigations, Great Britain Ministry of Agriculture, Fisheries and Food. H.M. Stationery Office, London, 1969, 1-38.
 22. Shafi M, Quddus MMA, Nahar S. Observations on the biology of the prawns *Macrobrachium lamarrei* Edwards and *Macrobrachium mirabilis* Kemp. *Bangladesh Journal of Scientific and Industrial Research*, 1977; 12(3-4):137-142.
 23. Tzomos Th, Koukouras A. Redescription of *Palaemon antennarius* H. Milne Edwards, 1837 and *Palaemon migratorius* (Heller, 1831) (Crustacea, Decapoda, Palaemonidae) and description of two new species of the genus from the circum-Mediterranean area. *Zootaxa*, 2015; 3905(1):27-51.
 24. Vaslet A, Bouchon-Navaro Y, Louis M, Bouchon C. Weight-length relationships for 20 fish species collected in the mangroves of Guadeloupe (Lesser Antilles). *Journal of Applied Ichthyology*, 2008; 24(1):99-100.
 25. Wickins JF. Experiments on the culture of the spot prawn *Pandalus platyceros* (Brandt) and the giant freshwater prawn *Macrobrachium rosenbergii* (deMan). 1972; *Fishery Investigation London Series*. 1972; 2:27(5):1-23.
 26. Willführ-Nast J, Rosenthal H, Udo PJ, Nast F. Laboratory cultivation and experimental studies of salinity effects on larval development in the African River prawn *Macrobrachium vollenhovenii* (Decapoda, Palaemonidae). *Aquatic Living Resources*, 1993; 6:115-137.