



International Journal of Fisheries and Aquatic Studies

E-ISSN: 2347-5129

P-ISSN: 2394-0506

(ICV-Poland) Impact Value: 5.62

(GIF) Impact Factor: 0.549

IJFAS 2021; 9(1): 135-141

© 2021 IJFAS

www.fisheriesjournal.com

Received: 13-11-2020

Accepted: 19-12-2020

Gyanendra Bahadur Chand

Associate Professor, Aquatic
Toxicology Lab., Department of
Zoology, Patna University,
Patna, Bihar, India

Khushbu Kumari

Research Scholar, Department of
Zoology, Patna University,
Patna, Bihar, India

Aakanchha

Research Scholar, Department of
Biotechnology, Patna
University, Patna, Bihar, India

Corresponding Author:

Khushbu Kumari

Research Scholar, Department of
Zoology, Patna University,
Patna, Bihar, India

A comparative study of the hematological parameters of *Clarias batrachus* (Linnaeus, 1758) and *Clarias gariiepinus* (Burchell, 1822) from North Bihar, India

Gyanendra Bahadur Chand, Khushbu Kumari and Aakanchha

DOI: <https://doi.org/10.22271/fish.2021.v9.i1b.2395>

Abstract

The genus *Clarias* (Family: *Clariidae*; order: *Siluriformes*) incorporates air breathing Clariid catfishes, having ability to survive out of water for longer duration. *Clarias batrachus* (Linnaeus, 1758) [24], commonly called as 'Magur', is an endogenous species of India and is very popular in Asiatic countries because of its taste and medicinal properties. *Clarias gariiepinus* (Burchell, 1822), an African catfish commonly called as 'Thai magur, is deadly carnivorous exotic species introduced to India through various trade means. This exotic species has undergone tremendous adaptation to the physicochemical condition of the aquatic bodies of Bihar, India, resulting in a considerable decline of the native species *C. batrachus* from the local fish market. The aim of the present study was to investigate and establish reference ranges by comparing haematological indices of two closely allied fresh water air breathing species of family *Clariidae* i.e. *C. batrachus* and *C. gariiepinus* from the wetland of North Bihar. In the present study hematological indices like red blood corpuscles (RBC), total and differential white blood corpuscles (WBC), hematocrit (Ht), haemoglobin (Hb), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC) were measured in the blood samples of both the fish species using standard methods. The values were determined as mean \pm SEM for thirty (n=30) replicates in each parameter. One-way analysis of variance (ANOVA) was used to compare the haematological findings between and within these two species of catfishes. Most of the haematological parameters differ non-significantly ($p < 0.05$) between *C. batrachus* and *C. gariiepinus* except haemoglobin and leucocytes, which showed highly significant differences. The findings of the present study will give an insight of the actual reference range of various haematological parameters and interspecific variations between these two species with regard to the blood profile. Furthermore, these indices will provide reliable information on the metabolic disorders, deficiencies and chronic stress status of the fishes before they become apparent in a clinical setting. It will definitely help in raising the status of aquaculture in state of Bihar.

Keywords: Blood, *Clarias batrachus*, *Clarias gariiepinus*, Haematological analysis, North Bihar, India

Introduction

Health and physiological status of the fish can be assessed effectively by considering haematological indices as bio-indicators (Clauss *et al.*, 2008; Adeyemo *et al.*, 2009; Siwicki *et al.*, 1994) [14, 2, 52]. In addition, these parameters play a significant role in the diagnosis of disease and assessment of metabolism of fishes living in different ecological environments by reflecting the correct changes within them (Clarence and Hickey, 1982; Cengizler and Şahan, 2000) [13, 12]. The exhaustive studies made on the haematological indices have successfully been used to establish the phylogenetic relationship among certain species of fishes (Cameron 1970; Larsson *et al.* 1976; Atkinson and Judd 1978; Putnam and Freel, 1978; Filho *et al.* 1992) [10, 33, 5, 45, 92]. The blood in fishes measures 1.3-7% of their total body weight. It ensures gas exchange between the organism and the environment. It plays a detrimental role in accomplishment of different metabolic processes. It facilitates the transportation of all kinds of nutrients, enzymes, hormones, neurotransmitters and all metabolic intermediaries in the different parts of body, apart from helping in transporting nitrogenous wastes outside the body of the fish. The functional status of the oxygen carrying capacity of the blood stream in fish can be effectively assessed by several hematological indices such as hematocrit (Hct), hemoglobin (HB), red blood cells (RBCs) *etc.* Hence these indices are convincingly used as bio-indicators of the patho-physiological condition or sub-lethal stress response in fish to

various endogenous or exogenous changes in the aquatic environment (Cataldi *et al.* 1998; Belanger *et al.* 2001; Shah and Altındağ, 2004a; Adhikari *et al.*, 2004; Radu *et al.*, 2009; Smith, 1945) [11, 7, 49, 3, 46, 53].

Fishes being poikilothermic vertebrates, experiences considerable changes in the metabolic contents of the blood in response to change in various environmental factors *viz.* bacteria, parasites, water temperature, oxygen content, pH *etc.* (Bullis, 1993) [9]. Besides, the blood components exhibit a wide variety of genetic and physiological variations. The aquatic biotope, fish species, age, sexual maturity and health status influence the changes in haematological parameters (Blaxhall, 1972; Patriche *et al.* 2011; Radu *et al.* 2009) [42, 46]. An ideal reference range of various blood components is considered as reliable descriptors of healthy fish under natural conditions (Cataldi *et al.*, 1998) [11]. Any kind of evaluation of haematological parameters is based on the normal reference range under ideal natural condition.

Clarias batrachus, commonly known as ‘magur’ is an endogenous Asian species, which is commercial highly significant in India and other Asian countries including Bangladesh, Thailand, Vietnam, Malaysia and Indonesia due to its excellent nutritional status with high Vitamin D content, low level of omega-3 fatty acids and a much higher proportion of omega-6 fatty acids as well as its high medicinal and therapeutic value (Mollah & Karim, 1990; Hossain *et al.*, 2006; Islam *et al.*, 2007) [39, 24, 27]. After being originated in India, it spreads globally (Kottelat, 2001; Lever, 1996) [32, 34]. Four sub species of *Clarias batrachus* are known to be prevalent in South East Asia, Java and India as *C. batrachus*; *C. aff. batrachus* “Indochina”; *C. aff. batrachus* “Sundaland” and *Clarias magur* (Hamilton, 1822) (Ng & Kottelat, 2008) [40]. The invasion of a morphologically alike and deadly carnivores African species *Clarias gariepinus* (Burchell, 1822) in India and their ability to hybridize naturally with the native species and excessive predation on them have potentially threaten the survival of *Clarias batrachus* (Thakur, 1998; Khedkar *et al.*, 2010) [60, 28]. The culture and import of *C. gariepinus* in India was totally banned in 1997 by the Ministry of agriculture in India (Gopi & Radhakrishnan, 2002) [20]. But due to prolific breeding ability, excessive acclimatization in the Indian condition and simplicity in rearing, this exotic species has compelled the fishermen in Bihar and other states of India to culture it on large scale, setting aside the ban and restriction imposed by the ministry of agriculture, Government of India (Singh & Lakra, 2006).

The purpose of the present study was to characterize base-line concrete haematological indices of *Clarias batrachus* and *C. gariepinus*, and to establish a reference point that can be efficiently applied for monitoring of fish health, disease status, reproductive potentials and various eco-toxicological studies. The finding of the present study can be correlated with the advanced mitochondrial genome analysis of these two closely allied species of family *Clariidae*. It was assumed that a systemic study of their haematological indices may provide information about interspecies differences between them and adaptations of the exotic species to the aquatic conditions of Indian subcontinent.

Materials and methods

Biological sampling and acclimatization

A total of 60 adult specimens, 30 each of *C. batrachus* and *C. gariepinus* ranging from 60-110 gm ranging from 60-110 GM and size between 4.6”-7.1” were collected from Jalalgarh fish

farm, Purnia, Bihar, India during June 2019. Fishes were at first brought to laboratory and were disinfected with 0.1% KMnO₄ solution and kept in different sized large plexi glass aquaria (capacity 50L, 80 L and 100L) having DE chlorinated, aerated tap water at normal temperature and pressure. Fishes were acclimatized to the ideal laboratory condition for 15 days as per standard method of APHA (2005). Fishes were fed *ad libitum* and the fresh water was changed regularly in the morning hours of every day. To maintain normal temperature and pressure of water, cooler and exhaust were used around the aquarium.

Collection of blood sample

Fishes were cold anaesthetized using MS222 (Imanpoor *et al.*, 2010) [26]. The peripheral blood was collected by puncturing the posterior caudal vein with heparin treated 21 gauge x 0.5 inch needle fitted with 2 ml syringe as per the standard protocol of Schmitt *et al.* (1999). The collected blood samples were then transferred into microtubes, containing EDTA (ratio 1.26 mg/0.6 ml) as the anticoagulant agent and were gently mixed by carefully turning it upside down. The collection tubes were then kept in the sampling boxes containing ice. The time elapsing from capture to blood withdrawal was less than 5 min.

Haematological analysis

In the experimental protocol, estimation of Haemoglobin (HB) was done with a hemoglobin test kit (Diagnova, Ranbaxy, India) using the cyanmethemoglobin method (Blaxhall and Daisley, 1973; Tanyer, 1985) [8, 58]. “Microhematocrit Technique” was employed for determination of Hct (Blaxhall and Daisley, 1973; Konuk, 1981; Şahan and Cengizler, 2002) [8, 31, 47]. The total red blood cells (tRBCs) were counted using an improved Neubaur haemocytometer (Shah and Altındağ 2004a) [49]. Blood was diluted 1:200 with Hayem’s fluid (Mishra *et al.*, 1977) [38]. Erythrocytes were counted in the loaded haemocytometer chamber and total numbers were reported as 10⁶ mm⁻³ (Wintrobe, 1967) [63]. The erythrocyte indices were calculated according to the following formula: Mean Corpuscular Volume (MCV) (µ3/femtolitre) = HCT (%) ÷ RBC (10⁶/mm³) x 10; Mean Hemoglobin Concentration (MCH) (PG/picogram) = HB (g/100 mL) ÷ RBC (10⁶/mm³) x 10; Mean Corpuscular Hemoglobin Concentration (MCHC) (%) = HB (g/100mL) ÷ HCT (%) x 10 (Kocobatmaz and Ekingen, 1984) [29].

Total count of WBC Total white blood cells (WBC) were counted using an improved Neubaur haemocytometer (Shah and Altındağ 2005; Mgbenka *et al.*, 2003) [50, 37]. Blood was diluted 1:20 with Turk’s diluting fluid and placed in haemocytometer. 4 large (1 SQ mm) corner squares of the haemocytometer were counted under the microscope (Olympus) at 640 X. The total number of WBC was calculated in mm³ x 10³ (Wintrobe, 1967) [63].

Statistical analysis

Results are presented as mean with standard error of mean (SEM) of thirty replicates in each case. The results were subjected to student’s t-test. One Way analysis of Variance (ANOVA) was used to determine significant differences in between experimental groups and significance levels of these differences of all parameters measured (Hayran and Özdemir, 1995) [23]. The level of significance was considered at *p* < 0.05. SPSS 10.0 software was used for statistical analyses.

Results

The haematological value obtained for the two species *i.e.* *Clarias batrachus* (Linnaeus, 1758) [24] and *Clarias*


gariepinus (Burchell, 1822) have been presented as Mean ± SEM in each case in Table – I.

Table 1: Hematological parameters of *Clarias batrachus* (Linnaeus, 1758) [24] and *Clarias gariepinus* (Burchell, 1822)


Demonstrated parameter	Sample size	<i>Clarias batrachus</i>		<i>Clarias gariepinus</i>		Calculated F value	Table F value
		Mean	SEM	Mean	SEM		
RBC (x10 ⁶ / μl)	30	2.726	0.168	2.40	0.362	0.803	5.31
PCV (%)	30	37	5.44	34.26	3.85	0.168	5.31
Hb (g/dl)	30	9.29	0.82	6.52	0.26	10.25*	5.31
MCV(fl)	30	138.6	22.23	116.2	11.3	0.811	5.31
MCH(PG)	30	34.48	3.33	35.37	5.44	0.019	5.31
MCHC (g/dl)	30	29.01	2.87	29.98	2.65	0.061	5.31
WBC (x10 ³ / μl)	30	106.56	23.41	70.82	16.68	1.545	5.31
Lymphocyte (%)	30	64.56	4.27	49.22	6.46	3.917	5.31
Monocyte (%)	30	1.6	0.40	3.0	0.31	7.53*	5.31
Eosinophils (%)	30	0.60	0.39	0.80	0.67	0.13	5.31

Figures in each parameters have been shown as Mean ± SEM, the sample size in each in case is n = 30, * significant at p <


0.05. One way analysis of Variance was done. Calculated value has been compared with table value.



Graph 1



Graph 2



Graph 3

Discussions

The blood is considered as a mirror in which, majority of the vital processes of an organism taking place, are reflected. The blood of teleost efficiently reflects the essence of the metabolic processes in fishes (Hawkins, 1971) [22]. Besides, toxicological research and environmental monitoring studies have consistently included fish blood as a possible indicator of patho-physiological changes in fishes. Therefore, it is incorporated as essential component of the fisheries management in India (Gupta and Gupta, 1981; Sancho *et al.*, 2000; Barcellos *et al.*, 2003) [21, 48, 6]. The present study was aimed to establish a correlation among two morphologically identical closely allied species of catfishes *i.e.* *Clarias batrachus* (Linnaeus, 1758) [24] and *C. gariepinus* (Burchell, 1822) of family-*Clariidae* and order-*Siluriformes*, if any, with special reference to haematological parameters. The present study also deals about variations among blood parameters in these two closely related species. Functional status of the oxygen carrying capacity of the bloodstream in fishes can be assessed by a number of hematological indices such as haematocrit (HCT), hemoglobin (HB), Red blood cells (RBCs), MCH, MCHC, MCV, *etc.*

RBCs are the most numerous (98-99%) blood cells in fish (Fange 1994). In the present study, the level of RBCs in control *C. batrachus* and *C. gariepinus* were recorded as $2.72 \pm 0.168 \times 10^6/\mu\text{l}$ and $2.40 \pm 0.36 \times 10^6/\mu\text{l}$ respectively with an average range of $2.40\text{-}2.80 \times 10^6/\mu\text{l}$. The calculated F value is 0.803, which is much lower than the table value. Hence the difference in the RBC count in both the species are not significant at $p < 0.05$. However a slight difference in RBC count in both the species may be correlated to the difference in the size and swimming activities (Soldatov, 2005) [54]. Maheshwaran *et al.*, (2008) [36] has determined the total

erythrocytes count as $177 \pm 0.014 \times 10^6/\mu\text{l}$. Adedeji and adegbile (2011) [1] have reported total RBC count in *Clarias gariepinus* as $1.77 \pm 0.014 \times 10^6/\mu\text{l}$.

Fish erythrocytes are ellipsoidal and contain nucleus. However, erythrocyte count in fish is strongly dependent on environmental conditions, mostly on temperature and dissolved oxygen level. Therefore, the seasonal changes in erythrocyte numbers pose difficulties in establishing the hematologic reference values (Luskova 1997) [35]. In aquatic environment, O₂ availability may considerably vary, and fish react to hypoxic conditions with elevation of blood oxygen carrying capacity – a rapid release of stored cells from a splenic reservoir or an increase in erythropoietic rate (Houston, Roberts, and Kennington 1996) [25]. The metabolic rate in poikilotherm animals is greatly affected by ambient temperature which causes seasonal alterations in fish red blood cell parameters: fishes living in higher temperatures showed higher RBC (Sopińska 1983; Svetina *et al.* 2002) [55, 57]. Besides, water, pH, salinity affect erythrocyte count in fish.

RBC indices basically include MCV, MCH and MCHC, haemoglobin (HGB) and haematocrit (HCT) value. Mean cell (or corpuscular) volume (MCV) refers to the average size of the RBCs constituting the sample. Mean cell hemoglobin (MCH) represents to the average weight of hemoglobin in the RBCs in the sample. Mean cell hemoglobin concentration (MCHC) refers to the average concentration of hemoglobin in the RBCs contained within the sample. The saturation of an organism with oxygen is reflected by the concentration of haemoglobin and the organism itself tries to maintain them at homeostasis. A decrease in the erythrocyte count and in the percent of haematocrit indicates the worsening of an organism state and developing anaemia. The calculation of MCV, MCH

and MCHC were directly linked with reference to RBC, HCT and Hgb.

Erythrocytes of fish are sensitive to various environmental impacts but, according to Vosylienė (1999) [61], the basic quantitative red blood parameters (e.g. hematocrit, erythrocyte count or hemoglobin concentration) in fish tend to remain stable due to considerable compensatory potential. The damage of erythrocytes caused by adverse environmental factors may be compensated by the release of new cells from the spleen (erythrocyte storage organ) or/and head kidney (hematopoietic organ), which is indicated by increase in the frequency of immature cells in peripheral blood. In the present study, the level of MCV in control *C. batrachus* and *C. gariepinus* were recorded as 138.6 ± 22.23 FL and 116.2 ± 11.3 FL respectively with a range of 110-160 fl. The calculated F value is 0.811, which is much lower than the table value i.e. 5.31. It is non-significant ($p < 0.05$). However, the mean MCV for *Clarias gariepinus* from Nigeria have been reported as 99.575 ± 3.64 FL, ranging between 88-160.90 FL (Adedeji and Adegbile, 2011) [1].

The level of MCH in control *C. batrachus* and *C. gariepinus* were recorded as $34.48 \pm 3.33\%$ and $35.37 \pm 5.44\%$ respectively, with an average range of 30-42%. The calculated F value is 0.019, which is much lower than the table value. It is non-significant ($p < 0.05$). Although the MCH in *C. gariepinus* were earlier reported to be in range between 25.10-48.90pg with a mean value of 31.345 ± 1.20 pg (Adedeji and Adegbile, 2011) [1].

The level of MCHC in control *C. batrachus* and *C. gariepinus* been recorded as $29.01 \pm 2.87\%$ and $29.98 \pm 2.65\%$ respectively, with an average range of 26-32%. The calculated F value is 0.061, which is much lower than the table value. The difference in the concentration of MCHC in both the species are non-significant ($p < 0.05$). The MCHC in *C. gariepinus* was earlier reported to vary in between 26-32, with a mean value of 30.7 ± 0.41 (Adedeji and Adegbile, 2011) [1].

The packed cell volume (PCV) is a measurement of the proportion of blood that is made up of cells. The value is expressed as a percentage or fraction of cells in blood. It is one of the indicators of fish health and sometimes it is also called as hematocrit. The level of PCV % in control *C. batrachus* and *C. gariepinus* were recorded as 37 ± 5.44 and 34.26 ± 3.85 respectively, with an average range of 30-44%. The calculated F value is 0.168, which is much lower than the table value. It is non-significant ($p < 0.05$). Previously recorded mean PCV for *Clarias gariepinus* was $30 \pm 1.09\%$, with a range of 18-37% (Adedeji and Adegbile, 2011) [1].

The level of haemoglobin (g/dl) in control *C. batrachus* and *C. gariepinus* were recorded as 9.29 ± 0.82 and 6.52 ± 0.26 respectively. The calculated F value is 10.25, which is much higher than the table value. It is highly significant at $p < 0.05$. The higher concentration of haemoglobin in *C. batrachus* is linked with the high metabolic activities of the fish (Eisler, 1965) [65]. An abnormally high concentration of haemoglobin in *C. batrachus* (75.00 ± 4.04 g/dl) have been earlier reported (Maheshwaran, 2008) [36]. The mean haemoglobin concentration in *C. gariepinus* have been earlier reported as 9.28 ± 0.42 g/dl (Adedeji and Adegbile, 2011) [1].

White blood cells play a major role in the defense mechanism of the fish and consist of granulocytes, monocytes, lymphocytes and thrombocytes. Granulocytes and monocytes function as phagocytes to salvage debris from injured tissue and lymphocytes produce antibodies (Ellis *et al.*, 1978; Wedemeyer and Mcleay, 1981) [62]. The number of leucocytes

determines the extent of immune responses and the ability of the animal to fight infection (Douglass and Jane, 2010) [15]. Higher levels of WBC in any species are the indication of their efficient way of fighting with pathogen than other species. Fish thrombocytes are blood phagocytes that form a protective barrier (Tavares-Dias and Moraes, 2004; Prasad and Charles 2010; Prasad and Priyanka 2011) [59, 44, 43]. They express a large number of surface and intracellular molecules which are involved in immune function (Kollner 2004) [30]. They represent a link between innate and adaptive immunity (Passantino *et al.* 2005) [41] and participate in defense mechanisms efficiently (Stosik *et al.* 2001) [56]. In the present study, the level of WBCs in control *C. batrachus* and *C. gariepinus* were recorded as $106.56 \pm 23.41 \times 10^3/\mu\text{l}$ and $70.82 \pm 16.68 \times 10^3/\mu\text{l}$ respectively, with an average range of 60-115 $\times 10^3/\mu\text{l}$. The calculated F value is 1.54, which is much lower than the table value. It is non-significant ($p < 0.05$).

Lymphocytes are most common and variable leucocytes in healthy teleost. An increase in lymphocyte number may be the compensatory response of lymphoid tissues to the destruction of circulating lymphocytes (Shah and Altındağ, 2005) [50]. The level of lymphocyte in control *C. batrachus* and *C. gariepinus* were recorded as $64.56 \pm 4.27\%$ and $49.22 \pm 6.46\%$ respectively. The calculated F value is 3.91, which is much lower than the table value. It is non-significant at $p < 0.05$. The higher percentage of circulating lymphocytes in *C. batrachus* can be linked to its efficient humoral and cell mediated adaptive immune responses (Douglas and Jane, 2010) [15].

Monocytes in teleosts are the largest cell among different leucocytes, irregular in shape, having eccentric, large, heterochromatic nucleus and abundant slightly basophilic cytoplasm with vacuoles and other membrane bound organelles. In the present study, the concentration of monocyte (%) in control *C. batrachus* and *C. gariepinus* were recorded as 1.6 ± 0.40 and 3.0 ± 0.31 respectively. The calculated F value is 7.53, which is higher than the table value i.e. 5.31. It is significant at $p < 0.05$. True eosinophils are less common in teleost. The level of eosinophils (%) in control *C. batrachus* and *C. gariepinus* were recorded as 0.60 ± 0.39 and 0.80 ± 0.67 respectively. The calculated F value is 0.13, which is much lower than the table value. It is non-significant at $p < 0.05$. A concentration of comparatively higher eosinophils might be associated with parasitic eradication, allergy and chronic inflammation.

The present study summarizes some of the salient features of hematologic analysis in two closely allied species of air breathing fish *Clarias batrachus* (Linnaeus, 1758) [24] and *Clarias gariepinus* (Burchell, 1822) of family Clariidae, inhabiting the same aquatic bodies. The reference ranges obtained can be used later, as a sensitive index to monitor various and patho-physiological changes in fish due to xenobiotic stress. It will also help in monitoring both the health status of the fish and the pollution status of the aquatic bodies. Further studies are needed to trace the molecular and genetic basis of the significant variations in the haematological indices of these two closely allied fresh water air breathing fish species.

Acknowledgement

The authors are thankful to the Head, Department of Zoology, Patna University, Patna, Bihar for providing infrastructural facilities.

References

- Adedeji OB, Adejibile AFI. Comparative haematological parameters of the Bagrid catfish (*Chrysichthys nigrodigitatus*) and the African catfish (*Clarias gariepinus*) from Asejire Dam in southwestern Nigeria. *Journal of Applied Sciences and Research* 2011;7(7):1042-1046.
- Adeyemo OK, Okwilagwe OO *et al.* Comparative assessment of sodium EDTA and heparin as anticoagulants for the evaluation of haematological parameters in cultured and feral African catfish (*Clarias gariepinus*). *Brazilian Journal of Aquatic Science and Technology* 2009;13(1):19-24.
- Adhikari S, Sarkar B *et al.* Effects of cypermethrin and carbofuran on certain hematological parameters and prediction of their recovery in a freshwater teleost, *Labeo rohita* (Ham). *Ecotoxicol. Environ. Saf* 2004;58:220-226.
- APHA. Standard Methods for the Examination of Water and Wastewater. APHA, AWWA, WPCF, Washington 2005.
- Atkinson E, Judd FW. Comparative hematology of *Lepomis microlophus* and *Cichlasoma cyanoguttatum*. *Copeia* 1978;2:230-237.
- Barcellos LJG, Kreutz LC *et al.* Haematological and Biochemical characteristics of male jundia (*Rhamdia Quelen, Quoyand Gaima* RDT, Pimelodidae): changes after acute stress. *Aquacul. Res* 2003;34:1465-1469.
- Belanger JM, Son JH *et al.* Effects of short-term management stress and ACTH injections on plasma cortisol levels in cultured white sturgeon, *Acipenser transmontanus*. *Aquaculture* 2001;203:165-176.
- Blaxhall PC, Daisley KW. Routine haematological methods for use with fish blood. *Journal of Fish Biology* 1973;5(6):771-781.
- Bullis RA. Clinical pathology of temperate freshwater and Estuarine fish. In: Stoskopf MK (ed): *Fish Medicine*. Saunders Company, USA 1993, 232-239.
- Cameron JN. The influence of environmental variables on the hematology of pinfish (*Lagodon rhomboids*) and striped mullet (*Mugil cephalus*). *Comparative Biochemistry and Physiology* 1970;32:175-192.
- Cataldi E, Di Marco P *et al.* Serum parameters of Adriatic Sturgeon *Acipenser naccarii* (Pisces: *Acipenseriformes*): Effects of temperature and stress. *Comparative Biochemistry and Physiology* 1998;120:273-278.
- Cengizler İ, Şahan A. Seyhan Baraj Gölü ve Seyhan Nehrin de yaşayan aynalıazan (*Cyprinus carpio*, Linnaeus, 1758) larda bazı kan parametrelerinin belirlenmesi. *Turkish Journal of Veterinary and Animal Sciences* 2000;24:205-214.
- Clarence R, Hickey JR. Comparative hematology of wild and captive cunners. *Transactions of the American Fisheries Society* 1982;111:242-249.
- Clauss TM, Alistair DM *et al.* Hematologic disorders of fish. *Veterinary Clinics of North America: Exotic Animal Practice* 2008;11(3):445-462.
- Douglass JW, Jane KW (eds.) *Schalm's Veterinary Hematology*. John Wiley and Sons, Blackwell Publishing Ltd 2010, 1232.
- Eisler R. Erythrocytes count and Haemoglobin contents in nine species of marine teleosts Chesapeake Science 1965;6:119-120.
- Ellis EA, Roberts RJ *et al.* *Fish Pathology*. Bailliere Tindall, London 1978, 318.
- Fange R. Fish blood cells. In: Hoar Ws, Randall DJ, Farrel AP, editors. *Fish physiology*, vol.12 B. San Diego (CA): Academic Press Inc 1992-1994, 54.
- Filho DW, Elbe GJ *et al.* Comparative hematology in marine fish. *Comparative Biochemistry and Physiology* 1992;102:311-321.
- Gopi KC, Radhakrishnan C. Coastline habitat degradation and threat to the turtle nesting site at Kolavipalam beach, Khozhikode District, Kerala. *Environ Sci. Newsletter Zool. Surv. India* 2002;6:5-7.
- Gupta BK, Gupta RB. Haematological observations on the *Channa punctatus* (Bloch.) *Ind J Zool* 1981;9(2):51-55.
- Hawkins M, Thomasi T. *Fish Haematology Bibliography*. A J Fish Bi 1971;4:193-232.ol.
- Hayran M, Özdemir O. *Bilgisayar İstatistik ve Tıp. Hekimler Yayın Birliği*, Ankara 1995.
- Hossain QZ, Hossain MA *et al.* Artificial breeding and nursery practices of *Clarias batrachus* (Linnaeus,1758). *Sci. World* 2006;4:4.
- Houston AH, Roberts WC *et al.* Hematological Response in Fish: Pronephric and Splenic Involvements in the Goldfish, *Carassius auratus* L. *Fish Physiology and Biochemistry* 1996;15:481-489. <http://www.genomesize.com/cellsiz/fish.htm>.
- Imanpoor MR, Bagheri T *et al.* The anesthetic effects of clove essence in persian sturgeon, *Acipenser persicus*. *World Journal of Fish and Marine Sciences* 2010;2(1):29-36.
- Islam MN, Islam MS *et al.* Genetic structure of different population of walking cat fish (*C. batrachus* Linn.) in Bangladesh. *Biochem. Genet* 2007;45:647-62.
- Khedkar GD, Reddy AC *et al.* *C. batrachus* (Linn., 1758) Population is lacking genetic diversity in India. *Mol Biol Reports* 2010;37:1355-62.
- Kocabatmaz M, Ekingen G. Değişik Tür Balıklarda Kan Örneği Alınması ve Hematolojik Metotların Standardizasyonu. *Doğa Bilim Dergisi* 1984;8:149-159.
- Kollner B, Fischer U *et al.* Potential involvement of rainbow trout thrombocytes in immune functions: a study using a panel of monoclonal antibodies and RT-PCR. *Development and Comparative Immunology* 2004;28:1049-1062.
- Konuk T. *Pratik Fizyoloji Ders Kitabı*, Ankara Üniversitesi Veteriner Fakültesi Yayınları. Ankara Üniversitesi Basımevi. Ankara 1981.
- Kottelat M. *Fishes of Laos Colombo*. Edited by Colombo Wild life Heritage Trust Publication 2001.
- Larsson A, Johansson-Sjöbeck ML *et al.* Comparative study of some haematological and biochemical blood parameters in the fishes from Skagerrak. *Journal of Fish Biology* 1976;9:425-440.
- Lever C. *Naturalized fishes of the world*, London, England Academic Press 1996, 408.
- Luskova V. Annual cycles and normal values of hematological parameters in fishes. *Acta Sc. Nat. Brno* 1997;31(5):70-78.
- Maheswaran R, Devapaul A *et al.* Haematological studies of Freshwater fish, *Clarias batrachus* (L.) exposed to mercuric chloride. *International Journal of Integrative Biology* 2008;2(1):50-54.
- Mgbenka BO, Oluah NS *et al.* Effect of Gammalin 20 (Lindane) on differential white blood cell counts of the

- African catfish, *Clarias albopunctatus*. Bull of Environmental Contamination Toxicol 2003;71:248-254.
38. Mishra N, Pandey PK *et al.* Haematological parameters of an air-breathing mud eel, *Amphipnous cuchia* (Ham.). J Fish Biol 1977;10:567-573.
 39. Mollah MFA, Karim M. First record of Induced breeding of African Magur (*Clarias gariepinus*, Burchell) in Bangladesh. J Inland Soc Ind 1990;22:52-54.
 40. Ng HH, Kottelat M. The identity of *Clarias batrachus* (Linnaeus, 1758) with the designation of a neotype (Teleostei: Clariidae). Zool. J Linn Soc 2008;153:725-32.
 41. Passantino L, Cianciotta A *et al.* Do fish thrombocytes play an immunological role? Their cytoenzymatic profiles and function during an accidental piscine candidiasis in aquarium. Immunopharmacology and Immunotoxicology 2005;27:345-356.
 42. Patriche T, Patriche N *et al.* Serum biochemical parameters of farmed carp (*Cyprinus carpio*). Aquaculture, Aquarium, Conservation and Legislation – International Journal of the Bioflux Society 2011;4:137-140.
 43. Prasad G, Priyanka GL. Effect of fruit rind extract of *Garcinia gummi-gutta* on haematology and plasma biochemistry of catfish *Pangasianodon hypophthalmus*. Asian Journal of Biochemistry 2011;6:240-251.
 44. Prasad G, Charles S. Haematology and leucocyte enzyme cytochemistry of a threatened yellow catfish *Horabagrus brachysoma* (Gunther 1864). Fish Physiology and Biochemistry 2010;36:435-443.
 45. Putnam RW and, Freel RW. Hematological parameters of five species of marine fishes. Comparative Biochemistry and Physiology 1978;61:585-588.
 46. Radu D, Oprea L *et al.* Characteristics of haematological parameters for carp culture and Koi (*Cyprinus carpio* Linnaeus, 1758) reared in an intensive system. Bulletin UASVM, Journal of Animal Science and Biotechnology 2009;66:1-2.
 47. Şahan A, Cengizler İ. Seyhan Nehri (Adana kent içi bölgesi)’nde yaşayan benekli siraz (*Capoeta barroisi* Lorted, 1894) ve kızılğöz (*Rutilus rutilus*, Linnaeus, 1758)’de bazı hematolojik parametrelerin belirlenmesi. Turkish Journal of Veterinary and Animal Sciences 2002;26:849-858.
 48. Sancho E, Ceron JJ *et al.* Cholinesterase activity and haematological parameters as biomarkers of sublethal molinate exposure in *Anguilla anguilla*. Ecotoxicology Environment Safe 2000;46:81-86.
 49. Shah SL Altındağ A. Hematological parameters of tench (*Tinca tinca* L.) after acute and chronic exposure to lethal and sublethal mercury treatments. Bull. Environ. Contam. Toxicol 2004a;73:911-918.
 50. Shah SL, Altındağ A. Alterations in the immunological parameters of tench (*Tinca tinca* L.) after acute and chronic exposure to lethal and sublethal treatments with mercury, cadmium and lead. Turk J Vet. Anim Sci 2005;29:1163-1168.
 51. Singh AK, Lakra WS. Alien fish species in India: Impact and emerging scenario. J Ecophysiol Occup Health 2006;6:165-174.
 52. Siwick AK, Anderson DP *et al.* Dietary intake of immunostimulants by rainbow trout affects non-specific immunity and protection against furunculosis. Veterinary Immunology and Immunopathology 1994;41:125-139.
 53. Smith HM. The fresh water fishes of Siamor Thailand. Bulletin of US 439 National Museum (Smithsonian Institution) 1945;188:1-622.
 54. Soldatov AA. Peculiarities of Organization and Functioning of the Fish Red Blood System. Journal of Evolutionary Biochemistry and Physiology 2005;41:272-281.
 55. Sopińska A. Effects of Physiological Factors, Stress, and Disease on Haematological Parameters of Carp, with a Particular Reference to Leukocyte Pattern. I. Variability of Haematological Indices of Carp in Relation to Age and Gonad Maturity Stage. Acta Ichthyologica et Piscatoria 1983;13:59-81.
 56. Stosik H, Deptula W *et al.* Studies on the number and ingesting ability of thrombocytes in sick carps (*Cyprinus carpio*). Veterinarni Medicina 2001;46:12-16.
 57. Svetina A, Matasin Z *et al.* Haematology and Some Blood Chemical Parameters of Young Carp till the Age of Three Years. Acta Veterinaria Hungarica 2002;50:459-467.
 58. Tanyer G. Hematoloji ve Lobaratuvar Ders Kitabı. Ayyıldız AS, Ankara 1985.
 59. Tavares-Dias M, Moraes FR (eds.): Hematology of Teleost Fish. Villimpress, Ribeirao Preto, Sao Paulo 2004.
 60. Thakur NK. A biological profile of African cat fish *Clarias gariepinus* and impact of its introduction in Asia. In Ponniah A.G., Das P., Verma S.R. editors. Fish Genetics and Biodiversity Conservation, Muzaffarnagar (UP), India. Natcon Publication 1998;05:275-92.
 61. Vosylienė MZ. The Effect of Heavy Metals on Haematological Indices of Fish (Survey). Acta Zoologica Lituonica 1999;9(2):76-82.61.
 62. Wedemeyer GR, Mcleay DJ. Methods of determining the tolerance of fish to environmental stressors. In Pickering AD (ed) Stress and fish. Academic Press, New York, USA 1981, 247-275.
 63. Wintrobe MM. Clinical hematology 6th Eds. Lea and Febiger, Philadelphia, Library of Congress, Print USA 1967.