



International Journal of Fisheries and Aquatic Studies

E-ISSN: 2347-5129

P-ISSN: 2394-0506

(ICV-Poland) Impact Value: 5.62

(GIF) Impact Factor: 0.549

IJFAS 2021; 9(2): 234-240

© 2021 IJFAS

www.fisheriesjournal.com

Received: 01-01-2021

Accepted: 03-02-2021

S Marimuthu

Department of Zoology,
Annamalai University,
Annamalainagar, Chidambaram,
Tamil Nadu, India

S Puvaneswari

Professor, Department of
Zoology, Annamalai University,
Annamalainagar, Chidambaram,
Tamil Nadu, India

P Raja

PG Department of Zoology,
Hajee Karutha Rowther Howdia
College (Autonomous), Theni,
Tamil Nadu India

M Jiyavudeen

Department of Biochemistry,
TBML College, Porayar,
Nagapattinam, Tamil Nadu,
India

The variations in the nucleic acids DNA and RNA levels in fish *Mugil cephalus* inhabiting Uppanar estuary

S Marimuthu, S Puvaneswari, P Raja and M Jiyavudeen

DOI: <https://doi.org/10.22271/fish.2021.v9.i2d.2456>

Abstract

A survey was conducted in the Uppanar estuary to scrutiny the levels of DNA and RNA from the selected tissues of the test fish *Mugil cephalus*. The test fish was collected periodically from three different stations (station 1, station 2 and station 3) for a period of one year. In the present study, a significant decrease ($P < 0.05$) in the nucleic acid level, such as DNA and RNA was noticed in selected tissues such as brain, liver, kidney and gill of fish *M. cephalus* in the order of liver, gill, brain, kidney. Station wise as well as season wise results revealed that, the maximum decrease in tissue DNA and RNA level was observed only in S2 during summer followed by post monsoon, pre monsoon and minimum decrease during monsoon. This may be because of this studied area (station 2) is subjected to increased pollution stresses, originating mainly from SIPCOT industrial areas, human activities such as disposal of agricultural, municipal, domestic and other wastes in substantial quantities.

Keywords: Uppanar estuary, DNA and RNA levels, seasonal changes, *M. cephalus*

Introduction

Industrial effluent and municipality wastes are the main reason for unwanted changes in the quality of water and its inhabiting fish (Shakir *et al.*, 2013; Ravikiran and Kulkarni, 2015) [9, 7]. Pollutants can directly affect cellular organelles and nucleic acids that may lead to cell cycle modulation, and apoptosis (Wang and Shi, 2001) [14]. Nucleic acids such as DNA and RNA level indicate the degree of protein synthesis.

It is based on the truth that the DNA level per cell is stable within the same species, and the RNA is largely ribosomic and differs with the protein synthesis rate. Nucleic acids are considered as important biomarkers in aquaculture as well as in eco-toxicology assessment due to their role in controlling various activities of cells (Miliou *et al.*, 1998) [6].

There are several laboratory studies with reference to alterations in nucleic acid level after exposure to various pollutants (Anuradha Shukla and Shukla, 2012; Habib and Abou Shehata Samah, 2013; Shakir *et al.*, 2012) [17, 5, 10].

RNA plays a significant role in protein synthesis hence depletion in RNA contents also results in depletion in protein level (Tripathi *et al.*, 2002) [16]. The growth of organisms is based on the increase in protein level and increase in RNA-content or ribosomal activity (Buckley *et al.*, 1999). Nucleic acid indices can be used as an estimate of recent growth among fish showing different kinds of growth dynamics, as these indices are sensitive to both changes in length- and weight specific growth rates (Hame and Rickwood, 1990).

Any change in nucleic acid metabolism results in the reduction in the DNA content. Furthermore, inhibition of DNA synthesis, might affect both protein as well as amino acid levels by decreasing the level of RNA in protein synthesis machinery (Das and Mukherjee, 2003). Studies on the changes in nucleic acid level on fishes in polluted natural environment are scanty. Hence, the goal of the present study is to quantify the changes in the nucleic acids (DNA and RNA) in brain, liver, kidney and gill tissues of the fish *M. cephalus* collected from different stations of Uppanar estuary.

Materials and Methods

The estuary flowing in the SIPCOT area has divided into three sampling stations, approximately 4 km apart from each other. The sampling stations are located at

Corresponding Author:

S Puvaneswari

Professor, Department of
Zoology, Annamalai University,
Annamalainagar, Chidambaram,
Tamil Nadu, India

Poondiyankuppam (S1), Sangolikuppam (S2) and Kudikadu (S3) in the coastal area of Cuddalore. The test fish *M. cephalus* were collected periodically from three different stations. The fish was sacrificed and the desired organs (brain gill, liver and kidney) were dissected out for quantification of DNA and RNA.

The nucleic acid levels (DNA and RNA) were determined by using the following methods:

Estimation of DNA by- Diphenylamine method of Schneider (1940)^[8].

Estimation of RNA by- Orcinol method of Schneider (1940)^[8].

Results and Discussion

In the present investigation, analysis of nucleic acids was performed in the fish *M. cephalus* collected from different stations of Uppanar estuary in Cuddalore district. The results showed that the maximum decrease in DNA and RNA level was observed at S2 than S1 and S3 (Figs. 1-8). In confirmative with this result, Habib and Samah (2013)^[5] have also noticed a station wise change in the levels of nucleic acid and proteins in cat fish collected from different stations in Egypt. They have correlated these results with the accumulation of heavy metals Cd and Pb in the tissues of the test fish. This consistent report was also made by Ali and Shakoori (2011)^[1]. They have noticed a significant decrease in the levels of DNA and RNA in the test fish, *Tor putitora*

collected from a polluted site of River Kabul than the same fish collected from non polluted Warsak Dam. This maximum reduction in station 2 may be due to the higher level of discharges of SIPCOT industrial effluents with a number of known and unknown pollutants from various sources at this station (Balakrishanan, 2014)^[3]. The fishes involved in long term exposure may have resulted in failure of the body defence mechanism and could have caused tissue degeneration.

In the present study, the decrease in DNA and RNA level was found to be in the order of liver, gill, brain, kidney (Figs. 1-8). The variations in the levels of nucleic acids in tissue samples show the tissue specificity of the nucleic acids. The level of DNA in different tissue indicates cell numbers (WHO, 1984; Cui *et al.*, 1997)^[15, 4]. The fishes involved in long term exposure may have resulted in failure of the body defence mechanism and could have caused tissue degeneration. The reason for high reduction in DNA and RNA levels in selected tissues may be either due to the heavy metal toxicity or to the pesticide toxicity in *M. cephalus* and might have caused genotoxic action by decreased mitotic index and disturbed cell division (Topktas *et al.*, 1996)^[12] or due to inhibitory action of heavy metals on DNA and RNA synthesis (Walter *et al.* 1980)^[13] or by cell death due to focal necrosis (Shivanandappa and Krishnakumari, 1981)^[11]. The decrease in the nucleic acid levels may also be due to the increased activity of liver phosphatases.




Fig 1: DNA content (mg/gm) in Brain tissue of the fish *M. cephalus* collected from the chosen stations (1, 2 and 3) of Uppanar estuary during various seasons 3




Fig 2: DNA content (mg/gm) in Liver tissue of the fish *M. cephalus* collected from the chosen stations (1, 2 and 3) of Uppanar estuary during various seasons 3




Fig 3: DNA content (mg/gm) in Kidney tissue of the fish *M. cephalus* collected from the chosen stations (1, 2 and 3) of Uppanar estuary during various seasons 3




Fig 4: DNA content (mg/gm) in Gill tissue of the fish *M. cephalus* collected from the chosen stations (1, 2 and 3) of Uppanar estuary during various seasons




Fig 5: DNA content (mg/gm) in Brain tissue of the fish *M. cephalus* collected from the chosen stations (1, 2 and 3) of Uppanar estuary during various seasons




Fig 6: RNA content (mg/gm) in Liver tissue of the fish *M. cephalus* collected from the chosen stations (1, 2 and 3) of Uppanar estuary during various seasons 3




Fig 7: RNA content (mg/gm) in Kidney tissue of the fish *M. cephalus* collected from the chosen stations (1, 2 and 3) of Uppanar estuary during various seasons 3




Fig 8: RNA content (mg/gm) in Gill tissue of the fish *M. cephalus* collected from the chosen stations (1, 2 and 3) of Uppanar estuary during various seasons 3

Thus, Nucleic acids and protein contents are regarded as important biomarkers of the metabolic potential of cells, as these play the main role in regulating different activities of cells. Their ratios also provide significant information about the way in which, or the mechanism by which, these contents regulate the multifaceted activities of cells. Therefore, there is a necessary to develop and apply efficient technologies to notice, manage, and correct human induced degradation of aquatic systems.

Conclusions

Based on the results of the present study it can be concluded that the significant decrease in the DNA and RNA content might be due to the discharges of SIPCOT effluents with a number of known and unknown pollutants. The reasons for the decreased levels of nucleic acid are the enzyme inhibition, mitotic changes, impairment of nucleic acid metabolism, increased activity of phosphatases and the degradation of cells. On compilation of all these results it appears that the disruption of DNA synthesis might have reflected in RNA synthesis.

Acknowledgement

The authors would like to acknowledge the help and support stretched by the Department of Zoology, Annamalai University, Chidambaram in Tamil Nadu, India.

Reference

1. Ali MY, Shakoori AR. Hepatic Responses of A Freshwater Fish Against Aquatic Pollution Pakistan J Zool 2011;43(2):209-221.
2. Alaa H Al-Darraj. The actual truth about different procedures of DNA and its extractions processes. Int. J Adv Biochem Res. 2020;4(2):33-36. DOI: 10.33545/26174693.2020.v4.i2a.54
3. Blakrishnan K. Studies on the impact of industrial pollutants of Uppanar estuary on chosen tissues of Lates calcarifer, Ph.D. Thesis, Fac. Sci, Annamalai University 2014.
4. Cui L, Locatelli L, Xie MY, Sommadossi JP. Effect of nucleoside analogs on neurite regeneration and mitochondrial DNA synthesis in PC-12 cells, J Pharmacol. Exp 1997;280:1228-1234.
5. Habib AM, Samah AS. Effect of Heavy Metals Pollution on Protein iosynthesis in Catfish Journal of Water Resource and Protection 2013;5:555-562.
6. Miliou H, Zaboukas N, Moraitou-Apostolopoulou N. Biochemical composition, growth and survival of the Guppy, *Poecilia reticulata*, during chronic sublethal exposure to cadmium. Archives of Environmental Contamination and Toxicology 1998;35:58-63.
7. Ravikiran K, Kulkarni RS. Nucleic Acid Content in Male Fresh Water Fish *N. notopterus* Exposed to Copper Sulphate. International Letters of Natural Sciences Online: ISSN: 2300-9675, 2015;33:1-8.
8. Schneider WC. Academic press, New York 1940, 235-340.
9. Shakir HA, Chaudhry AS, Qazi JI. Impact of anthropogenic activities on physico-chemical parameters of water and mineral uptake in Catla catla from river Ravi, Pakistan. Environ Monit Assess 2013;185:2842-3833.
10. Shakir HA, Qazi JI, Shakoor AC. Examining muscles of *Cirrhinus mrigala* for biochemical parameters as a bio-indicator of water pollution by municipal and industrial effluents into River Ravi, Pakistan Int Aquat Res 2012-2014;6:221-228.
11. Shivanandappa T, Krishnakumari MK. Histochemical and biochemical changes in rats fed dietary Benzenehexachloride. Ind. J exptl. Boil 1981;19:1163-1168.
12. Topktas ME, Renci Zogullari, Ila HB. *In-Vivo* chromosomal aberrations in bone marrow cells of rats with marshal. Mut. Res 1996;371:259-264.
13. Walter Z, Czajkawska A, Lipecka K. Effects of malathion on genetic material of human lymphocyte stimulated by phytohemagglutinin (PHA). Hum. Genet 1980;53:375-381.

14. Wang S, Shi X. Molecular mechanisms of metal toxicity and carcinogenesis. Mol. Cell. Biochem 2001;222:3-9.
15. WHO (World Health Organization), Guide Lines for Drinking Water Quality, 3rd Edition, Geneva 1984.
16. Tripathi RD, Gupta DK, Rai UN, Inouhe M. Impacts of fly-ash on soil and plant responses. Journal of Plant Research 2002;115(6):401-409.
17. Anuradha Shukla, Shukla JP. Distillery effluent induced alterations in the nucleic acid and protein during testicular cycle of *Colisa fasciatus*, (Bl. & Schn), a tropical freshwater perch. International Journal of Pharma and Bio Sciences 2012, 3(1).