



International Journal of Fisheries and Aquatic Studies

E-ISSN: 2347-5129

P-ISSN: 2394-0506

(ICV-Poland) Impact Value: 76.37

(GIF) Impact Factor: 0.549

IJFAS 2023; 11(4): 32-36

© 2023 IJFAS

www.fisheriesjournal.com

Received: 19-04-2023

Accepted: 22-05-2023

Sandeep Shukla

P.G. Department of Zoology,
B.S.N.V.P.G. College, Lucknow,
Uttar Pradesh, India

Shareef Ahmad

P.G. Department of Zoology,
B.S.N.V.P.G. College, Lucknow,
Uttar Pradesh, India

Samiksha Lodhi

P.G. Department of Zoology,
B.S.N.V.P.G. College, Lucknow,
Uttar Pradesh, India

Aman Ahmad

P.G. Department of Zoology,
B.S.N.V.P.G. College, Lucknow,
Uttar Pradesh, India

Richa Shukla

Department of Zoology, Navyug
Kanya P.G. College, Lucknow,
Uttar Pradesh, India

Sanjive Shukla

P.G. Department of Zoology,
B.S.N.V.P.G. College, Lucknow,
Uttar Pradesh, India

Corresponding Author:

Sandeep Shukla

P.G. Department of Zoology,
B.S.N.V.P.G. College, Lucknow,
Uttar Pradesh, India

Background colour induced modification in general body coloration of freshwater prawn *Macrobrachium lamarrei* (Crustacea-Decapoda)

**Sandeep Shukla, Shareef Ahmad, Samiksha Lodhi, Aman Ahmad, Richa
Shukla and Sanjive Shukla**

DOI: <https://doi.org/10.22271/fish.2023.v11.i4a.2822>

Abstract

Colour changing in animals of different taxa is a vital phenomenon, which can also be exploited in terms of increasing demand of market therefore can prove economical. Freshwater prawn, *Macrobrachium lamarrei* (Crustacea-Decapoda) was exposed to five different background colours viz. white, red, green, blue and black along with one control i.e. translucent. Chromatophore density and type was found variable in different background colours accordingly. The number of chromatophores was found significantly increased in darker backgrounds in comparison to control and maximum number was recorded in black background. The order of darkness of prawns was observed as white<red<green<blue<black. Underlying mechanism of colour change and market economy perspective of darker prawns have been discussed.

Keywords: Pangasius, hybrid, digestive system, histology

1. Introduction

To cope with increasing demand of food for increasing human population, aquaculture is most promising and the major hope for future to provide better nutrition specially for developing countries like India. Among aquaculture products other than fishes, crustaceans, specially prawns are a future hope (Janaki Ram *et al.*, 2003; Jayasankar 2018; FAO, 2019) [9, 10, 6].

Crustaceans includes crabs, lobster, prawns etc, which are relished worldwide. Freshwater prawn aquaculture is a recent one and is more beneficial than marine prawn aquaculture. Fresh water prawn *Macrobrachium lamarrei* (Crustacea-Decapoda) is a smaller prawn, locally available throughout the year, having complete development in freshwater and promising candidate for freshwater aquaculture (Shukla and Sharma, 2010; Shukla *et al.*, 2017; Ahmad *et al.*, 2021; Sharma *et al.*, 2022) [26, 30, 1, 25]. This prawn species is also very important in terms of bio-indicator for environmental monitoring (Sharma and Shukla, 1990; Sharma and Shukla, 2006; Lodhi *et al.*, 2006; Lodhi *et al.*, 2004; Lodhi *et al.*, 2008; Lodhi *et al.*, 2009; Shukla and Sharma, 2010; Verma *et al.*, 2010; Ahmed *et al.*, 2021) [39, 23, 14, 15, 17, 16, 26, 35, 1]. Market value of prawns depends upon their colour because dark colour prawns are always preferred. By using suitable background colour the body colouration of cultured prawn can be managed and in this way market value can be enhanced (Parisenti *et al.*, 2011; Latscha 1989; Tume *et al.*, 2009; Shukla *et al.*, 2017; Palomera *et al.*, 2018) [20, 13, 34, 30, 19]. Considering the above facts present work is considered to observe the effect of background colouration on fresh water prawn, *Macrobrachium lamarrei* (Crustacea-Decapoda), a potential but “untapped” resource for fresh water aquaculture.

2. Material and Methods

Fresh water prawns, *Macrobrachium lamarrei* (Crustacea-Decapoda) (H. Milne Edwards) were collected from river Gomti, Lucknow (U. P.) – India, with the help of local fisherman and brought to the laboratory (N- 26° 49' 5" E- 80° 55' 58") in large plastic containers. Adult inter-moult staged *M. lamarrei* (Avg. length – 48.6±5.5 mm; weight – 1.107 ±0.26gm.) were

being utilized in experiments after 5-7 days acclimation to laboratory conditions. The animals were maintained in different background coloured containers (*viz.* White, blue, green, red and white) along with one control (translucent) were used to determine effect on coloration of prawn, containing dechlorinated water having physico-chemical characteristics as follows: - pH -7.66 ± 0.27 , Temperature $-28 \pm 1^{\circ}\text{C}$, D.O. -6.6 ± 0.74 mg/l, Total Hardness -268 ± 2.67 mg/l, Alkalinity -425 ± 11.36 mg/l (Sharma & Shukla, 1990; APHA *et al.*, 1998) [26, 2]. For experiment 15 animals in each tank were kept under normal photoperiod and controlled laboratory conditions like continuous aeration and feeding on alternate days along with control.

After 30 days coloration of animals of each container

were observed with naked eyes as well as with stereoscopic dissecting microscope and photographed. In present study no animal were sacrificed. All the observations were made on live animals. Experiment was replicated thrice.

4. Results and Discussion

Background colour of tank significantly affects prawn colouration. Density, type, distribution and structure of chromatophores vary according to the colour of background. It was observed that density of chromatophores increased in prawns of darker containers in comparison to lighter colour containers (Plate: 1-Fig. 1-12 & Plate: 2- Fig. 13-24).

The result shows that prawns are capable to modify their body colouration according to the background colour of experimental tanks. In red, green, blue and black colour backgrounds number of chromatophores are found to be increased while in white background number of chromatophores decreased than control (Plate:1-Fig. 1-12 & Plate:2-Fig. 13-24). Order of darkness in prawns according to background coloration is as follows: white<Red<Green<Blue<Black.

Colouration is a vital phenomenon of an organism as it plays crucial role in biology of organism. Change in pigmentation aid poikilotherms in visual communication, courtship, and survival. Chromatophores of crustacean have been assigned for various functions like photoprotection, cryptosis and/or thermoregulation (Fuhrman, 2011; Shukla *et al.*, 2017) [8, 28]. The pigment can be either dispersed throughout the cell, which gives a dark appearance, or it can be aggregated around the nucleus, which gives a pale appearance. Background colouration also affects several vital activities of prawn including metamorphosis and survival. It was reported that darker background enhances survival of larvae of *M.*

resenbergii (Sebastian and George, 1994; Yasharian *et al.* 2005; Shelke, 2010; Borisov *et al.*, 2022) [22, 36, 27, 5].

Chromatophores of prawns are present in hypodermal layer between the exoskeleton and abdominal muscle. Colour change in crustacean may be either due to physiological or due to morphological mechanisms or both. Physiological mechanisms that influence prawn colour include carotenoid availability in the diet, background substrate colour, photoperiod, light intensity and temperature. Light intensity affects neuro-endocrine system present in eye stalk which via various hormones affects number of various chromatophores and their distribution.

The green/gray colour observed in prawn is due to the accumulation of crustacyanin, a protein-astaxanthin complex that becomes orange with complex dissociation on cooking etc. Astaxanthine is a chemical which is responsible for the colouration in crustaceans (Boonyaratpalin *et al.*, 2001; Menasveta *et al.*, 2004; Laohavisuti and Ruangdej, 2014) [4, 40, 12]. It can be either synthesized or obtained from the diet. Manipulation of diet with astaxanthine compound also makes colour darker due to presence of high amount of astaxanthine (Yamada *et al.*, 1990; Boonyaratpalin *et al.*, 2001; Arredondo Figueroa *et al.*, 2003; Tume *et al.*, 2009; Parisenti *et al.*, 2011; Wade *et al.*, 2012) [37, 4, 3, 34, 20, 36].

Colour pattern of prawn are also dependent on physico-chemical properties of water like pH, hardness, temperature etc. (Shukla *et al.*, 2017) [30]. Central accumulation as well as spreading of pigment in chromatophores is also responsible for colour change under influence of various physico-chemical conditions involving Ca^{++} influx (Fingerman, 1965; Reibero & McNamara, 2006, Palomera *et al.*, 2018) [7, 21, 19]. Observations of present study are in accordance of above studies. Darker prawns give bright orange colour after cooking which signifies freshness and relishes all over the world with higher price in market.

Resent study showed that it is possible to improve the overall appearance of prawn (raw and cooked) by manipulating background colour about the time of harvesting. These small freshwater prawns can be a good source of nutrient rich food along with good source of income and entrepreneurship. Size, weight, shape, and colour are the decisive parameters that determine the value of a fish, prawn or an organism in the market. Change in production streatgy using suitable culture techniques can prove beneficial. Use of the background colour manipulation may be a lower-cost alternative to intensify the colour of the prawn being cultivated which are not fed with food containing carotenoids or pigments.






Plate 1: Explanation of figures: Photomicrographs of freshwater prawn, *Macrobrachium lamarrei* Plate:1 Fig. 1-4: Control; Fig. 5-8: White Background; Fig. 9-12: Red Background; RS=Rostrum; EY=Eye; CR=Carapace; AB=Abdomen; CH=Chromatophore; TL=Telson; UR=Uropode






Plate 2: Explanation of figures: Photomicrographs of freshwater prawn, *Macrobrachium lamarrei* Plate: 2 Fig. 13-16: Green Background; Fig. 17-20: Blue Background; Fig 21-24: Blue Background

Conclusion

Outcome of present study indicates that manipulation of rearing tank colour changes prawns in darker colour which will increase its market price and can be used widely in freshwater prawn aquaculture. This technique is cost effective than costly diet manipulation for the same purpose. This technique is unique in terms of food value because pigmentation of prawns can be enhanced without any chemical treatment thus keeping the nutritional value intact and unchanged. This technique is effective, cheap and easy for freshwater prawn farmers so likely to adaptable.

Acknowledgements

Authors are thankful to the Prof. U. D. Sharma, Rtd. Professor, Department of Zoology, University of Lucknow, Lucknow for valuable suggestions, supervision and blessings, and also thankful to Head, P.G. Department of Zoology, B.S.N.V.P.G. College, Lucknow (U.P.) India for providing necessary lab facilities.

References

- Ahmed A, Lodhi S, Shukla S. Observations on feeding behaviour of fresh water prawn *Macrobrachium lamarrei* (Crustacea: Decapoda) Int. J. of Fish. and Aqua. Stud. 2021;9(6):109-112
- APHA. Standard Methods for the examination of water and waste waters. 20thEdn. APHA, AWWA and WPCF Washington; c1998.
- Arredondo-Figueroa JL, Pedroza-Islas R, Ponce-Palafox JT, Vernon-Carter EJ. Pigmentation of Pacific white shrimp (*Litopenaeus vannamei*, Boone, 1931) with esterified and saponified carotenoids from red chili (*Capsicum annum*) in comparison to astaxanthin. Revista Mexicana de Ingeniería Química, Iztapalapa. 2003;2(1):101-108.
- Boonyaratpalin M, Thongrod S, Supamattaya K, Britton G, Schlipalius LE. Effects of β -carotene source, *Dunaliella salina*, and astaxanthin on pigmentation, growth, survival and health of *Penaeus mondon*. Aquacult Res, Oxford. 2001;32:182S-190S.
- Borisov RR, Nikonova IN, Parshin-Chudin AV, Kovacheva NP. Effect of Background Color on the Coloration of Juvenile Giant Freshwater Prawn *Macrobrachium rosenbergii*. Russ. J of Ecol. 2022;53(1):40-47.
- FAO. The State of the World's Aquatic Genetic Resources for Food and Agriculture. FAO Commission on Genetic Resources for Food and Agriculture assessments. Rome; c2019.
- Fingerman M. Chromatophores. Physiol. Rev. 1965;45(2):296-339.
- Fuhrmann MM, Nygård H, Krapp RH, Berge J, Werner I. The adaptive significance of chromatophores in the Arctic under-ice amphipod *Apherusa glacialis*. Polar Biology. 2011 Jun;34:823-32.
- Janaki Ram K, Pandey AK. Status and prospects of aquaculture development with special reference to Uttar Pradesh. Brain Storming Session on Production Status and Potential of Fisheries in UP: Constraints and Opportunities. 2003, 24-38.
- Jayasankar P. Present status of freshwater aquaculture in India- A review. Indian Journal of Fisheries. 2018;65(4):157-165.
- Kalpna P, Meena P. Role of appendages in feeding behaviour in freshwater crab *Barytelphusa cunicularis* and prawn *Macrobrachium kistnensis* (Decapoda-Crustacea) Int. J. Rec. Sci. Res 2015;7(3):9327-9330.
- Laohavisuti N, Ruangdej U. Effect of Dietary Astaxanthin and Background Colour on Pigmentation and Growth of Red Cherry Shrimp, Neocaridina heteropoda. Kasetsart University Fish. Res. Bull. 2014;38(1) 1-7
- Latscha, T. The role of astaxanthin in shrimp pigmentation. Advances in Tropical Aquaculture, Advances in Tropical Aquaculture, Tahiti. 1989;9:319-325.
- Lodhi HS, Khan MA, Verma RS, Sharma UD. Acute toxicity of Copper Sulphonate to freshwater prawns. J. Environ. Biol. 2006;27(3):585-588.
- Lodhi HS, Khan MA, Verma RS, Sharma UD. Seasonal fluctuations in the total haemocyte counts (THC's) of freshwater prawn, Him. J Env. Zool. 2004;18(1):1-5.
- Lodhi HS, Shukla S, Sharma UD. Studies on structure of heart of two freshwater prawns under light microscope. Uttar Pradesh J. Zool. 2009;29(1):1-10.
- Lodhi HS, Tiwari KJ, Shukla S, Sharma UD. Copper induced fluctuations in total haemocyte counts (THCs) of freshwater prawn, *Macrobrachium lamarrei* (Crustacea-Decapoda). J. Env. Bio. Sci. 2008;22(2):135-142.
- Gangbé L, Dessouassi E, Houndonougbo PK, Brahim AA, Djimènou D, Agadjihouédé H, et al. Trophic regime of the giant freshwater shrimp *Macrobrachium vollohoveni* (Herklots, 1857) in the lower Oueme valley in Southern Benin. Int. J Biol. Sci. 2022;4(1):37-45. DOI: 10.33545/26649926.2022.v4.i1a.52
- Palomera MAA, Villasante FV, Martínez CM, González AM, Zapata DB. Background colour effect on the pigmentation of prawn *Macrobrachium tenellum*. Lat. Am. J. Aquat. Res. 2018 Jul;46(3):610-614.
- Parisenti J, Beirao LH, Mourino JL, Vieira F, Buglione CC, Maraschim M. Effect of background color on shrimp pigmentation. Boletim do Instituto de Pesca. 2011 Jan 1;37(2):177-82.
- Ribeiro M, McNamara JC. Calcium movements during

- pigment aggregation in freshwater shrimp chromatophores. Pigment cell research. 2007 Feb;20(1):70-7.
22. Sebastian CD, George KC. Observation on hatchery seed production of *Macrobrachium rosenbergii* in Kerala. Freshwater prawn farming in India (eds. by Thakur, NK, Tewari R. and Josep). 1994, 85-91.
 23. Sharma UD, Shukla S. Acute toxicity of heavy metals and detergent to fresh water prawn *Macrobrachium lamarrei* (Crustacea-Decapoda). Him. J Env. Zool. 2006;20(1):1-6.
 24. Farhana S Ghory, Quddusi B Kazmi and Feroz A Siddiqui. First report of laboratory reared developmental stages of *Palaemon sewelli* (KEMP, 1925) (Crustacea: Caridea: Palaemonidae: Palaemonidae). Int. J Biol. Sci. 2021;3(2):38-44.
DOI: 10.33545/26649926.2021.v3.i2a.79
 25. Sharma M, Lodhi S, Ahmed A. Freshwater prawn aquaculture: Prospects in U.P. Int. J of Fish. and Aqu. Stud. 2022;10(4):146-151
 26. Shukla S, Sharma UD. Smaller fresh water prawns: Their aquaculture potential and suitability as good laboratory model. In: Bioresources for food security and rural livelihood. Kulkarni G.K. and Pandey P.N.(Eds) PP.189-204, Narendra Publications, Delhi; c2010.
 27. Shelke ST, Belsare SS, Indulkar ST. Effect of rearing container's colour on metamorphosis and survival of larvae of *Macrobrachium rosenbergii* (De Man, 1879). Asian Fisheries Science. 2010;23(1):25-34.
 28. Shukla S, Shukla R, Sharma UD. Effect of detergent, linear alkyl benzene sulphonate on gills of freshwater prawn, *Macrobrachium lamarrei*. Anusandhaan. 2013;1(1):31-34.
 29. Shukla S, Shukla R, Sharma UD. Histopathological effects of detergent, Linear Alkyl benzene sulphonate on hepatopancreas of freshwater prawn, *Macrobrachium lamarrei* (Crustacea-Decapoda). International Journal of Fisheries and Aquatic Studies. 2021;9(6):230-233.
 30. Shukla S, Shukla R, Shukla, S, Ahmad A, Mishra A. Effect of temperature on chromatophores of freshwater prawn, *Macrobrachium lamarrei* (Crustacea-Decapoda). Anusandhaan. 2017;5(1):80-84.
 31. Tewari KJ, Verma RS, Kasherwani D, Lodhi HS, Shukla Sanjive, Sharma UD. Lead induced behavioural anomalies in freshwater prawn, *Macrobrachium dayanum* (Crustacea-Decapoda). Environment & Ecology. 2007;25S(3A):802-804.
 32. Tiwari KJ, Lodhi HS, Mishra A, Ahmad A, Lodhi S, Shukla S. Effect of lead nitrate on haemocyte morphology of freshwater prawn, *Macrobrachium dayanum* (Crustacea-Decapoda) U. P. J. of Zool. 2022;43(13):80-87.
 33. Tiwari KJ, Lodhi HS, Shukla S, Sharma UD. Effects of Lead nitrate on Oxygen consumption of fresh water prawn, *Macrobrachium dayanum* (Crustacea-Decapoda); 2009; Environ. Conser. J 2009 Dec 21;10(3):9-13.
 34. Tume RK, Sikes AL, Tabrett S, Smith DM. Effect of background colour on the distribution of astaxanthin in black tiger prawn (*Penaeus monodon*): Effective method for improvement of cooked colour. Aquaculture. 2009 Nov 1;296(1-2):129-35.
 35. Verma DR, Lodhi HS, Tiwari KJ, Shukla S, Sharma UD. Copper sulphate induced changes in scaphognathite oscillations and oxygen consumption of fresh water prawn, *Macrobrachium lamarrei* (Crustacea-Decapoda). Journal of Applied and Natural Science. 2010 Jun 1;2(1):34-7.
 36. Wade NM, Anderson M, Sellars MJ, Tume RK, Preston NP, Glencross BD. Mechanisms of colour adaptation in the prawn *Penaeus monodon*. Journal of Experimental Biology. 2012 Jan 15;215(2):343-50.
 37. Yamada S, Tanaka Y, Sameshima M, Ito Y. Pigmentation of prawn (*Penaeus japonicus*) with carotenoids: I. Effect of dietary astaxanthin, β -carotene and canthaxanthin on pigmentation. Aquaculture. 1990 Jun 15;87(3-4):323-30.
 38. Yasharian D, Coyle SD, Tidwell JH, Stilwell WE. The effect of tank colouration on survival, metamorphosis rate, growth and time to metamorphosis freshwater prawn (*Macrobrachium rosenbergii*) rearing. Aquaculture Research. 2005 Feb;36(3):278-83.
 39. Sharma UD, Shukla S. Behavioural dysfunctions of fresh water prawn *M. lamarrei* (Crustacea-Decapoda) following exposure to a synthetic detergent, Linear Alkyl Benzene Sulphonate. Biological Memoirs. 1990;16(1-2):58-61.
 40. Menasveta P, Piyatiratitvokul S. Effect of different culture system on growth, survival, and production of the giant freshwater prawn. In: Giant Prawn Farming (ed. by New, M. B.). Elsevier Scientific Publishing Company, Amsterdam Oxford, New York. 1982;10:175-189.