



# International Journal of Fisheries and Aquatic Studies

ISSN: 2347-5129

(ICV-Poland) Impact Value: 5.62

(GIF) Impact Factor: 0.352

IJFAS 2015; 2(5): 363-365

© 2015 IJFAS

www.fisheriesjournal.com

Received: 19-04-2015

Accepted: 22-05-2015

**Ana Paula Madeira Di Beneditto**  
Universidade Estadual do Norte  
Fluminense Darcy Ribeiro,  
Centro de Biociências e  
Biotecnologia, Laboratório de  
Ciências Ambientais, Av. Alberto  
Lamego, 2000, Campos dos  
Goytacazes, RJ, 28013-602,  
Brazil.

## What drives the cannibalism of *Trichiurus lepturus* (Linnaeus, 1758) in the coastal area of southeastern Brazil (21-22°S)?

**Ana Paula Madeira Di Beneditto**

### Abstract

The present study was aimed to investigate the causes that drive the cannibalism of the teleost fish, *Trichiurus lepturus*, in the coastal area of southeastern Brazil (21-22°S) whereas the feeding behaviour was recorded through analyzing of the stomach contents. In this area, the population density of this predator fish is high, but there are many food resources available. Therefore, the highest cannibalism rate couldn't be expected. However, conspecifics intake is nutritionally more advantageous for the predator compared with other prey species, once they provide high protein content and calorific value. In this sense, the quality of food resources drives the cannibalism of this species locally.

**Keywords:** cannibalism; predator-prey interactions; southeastern Brazil; *Trichiurus lepturus*.

### 1. Introduction

Cannibalism is the act of a species eating the whole or major part of its own, irrespective of its stage of development. In teleost fish, cannibalism involves either at the egg or at post-hatching stages (larval, juvenile, or adult), comprising more than 10% of the recent fish families and fossil fish<sup>[13, 15]</sup>. In general, high cannibalism rate in fish is driven by: i) feeding habit, being more common in piscivores species; ii) shoaling behaviour; iii) size disparity among conspecifics, with larger specimens preying on small ones; iv) high fish density; and iv) nutritional state, with low food resources encouraging this feeding behaviour<sup>[15]</sup>. In general, cannibalism is a strategy to energy transfer from smaller to larger specimens, increasing in high population densities. By contrast, this feeding strategy would not be expected when other food resources are sufficient for the predators<sup>[1, 4, 11, 15, 16]</sup>.


*Trichiurus lepturus* (Linnaeus, 1758) is a teleost fish, known as large head hair tail, with economic importance as fishery resource around the world. The species forms shoals in brackish and marine waters, with aggregated feeding behaviour. There is size disparity among conspecifics, and significant diet shift was recorded during the ontogeny. Juvenile specimens are planktivores, while adult specimens are top predators (mainly piscivores), feeding on the most abundant prey available along its feeding sites<sup>[6, 10, 12]</sup>. Stomach contents analysis showed cannibalism as an important feeding behaviour for adult specimens<sup>[2, 12, 14]</sup>, but this is not a general rule along its distributional area<sup>[4, 5, 16]</sup>. Therefore, this study was planned for the first time the causes that drives the cannibalism of *T. lepturus* in the coastal area of southeastern Brazil (21-22°S) to understand its intraspecific trophic relationship.

### 2. Material and Methods

There are some scenarios to explain what drives the cannibalism behaviour of *T. lepturus* along its home range. The fish characteristics regarding feeding habits, shoaling behaviour, and size naturally favour cannibalism, but this feeding strategy is not a general rule for this species, as mentioned above. Then, one can say that a combination between population density and food resources drives the cannibalism, such as: i) high population density and many food resources = low cannibalism rate; ii) high population density and few food resources = high cannibalism rate; and iii) low population density and many food resources = low cannibalism rate. A scenario with low population density and few food resources might compromise the maintenance of a given *T. lepturus* population.

### Correspondence

**Ana Paula Madeira Di Beneditto**  
Universidade Estadual do Norte  
Fluminense Darcy Ribeiro,  
Centro de Biociências e  
Biotecnologia, Laboratório de  
Ciências Ambientais, Av. Alberto  
Lamego, 2000, Campos dos  
Goytacazes, RJ, 28013-602,  
Brazil.



**Fig 1:** Coastal area of southeastern Brazil where the cannibalism of *Trichiurus lepturus* was recorded (grey circle).

### 3. Results and Discussion

In southeastern Brazil, the cannibalism was recorded as the main feeding behavior of *T. lepturus* through stomach contents analysis [2, 3] (Fig 1). However, there is still no explanation to its high cannibalism rate. This fish is target of local commercial fisheries [9] and it is the main prey of the coastal dolphin *Sotalia guianensis* (van Bénédén, 1864) [7, 8], indicating its high density locally. Moreover, all prey of *T. lepturus* are abundant year round [2, 3, 10], and the low food resources was not a satisfactory explanation for its cannibal behaviour. The local scenario is high population density and many food resources. Therefore, the high cannibalism rate would not be expected.

In this area, the cannibalism of *T. lepturus* could be explained by the proximate-composition and caloric content provided by conspecifics. Juveniles' conspecifics, preferred prey of adults, have high protein content and caloric value [3]. Thus, their intake is nutritionally more advantageous for the predator compared with other prey, as small engrailed and clupeid fish species. Juveniles' conspecifics provide more protein for rapid growth and more energy to support wide movements along the continental shelf and reproductive demand.

### 4. Conclusions

In southeastern Brazil, between 21°S and 22°S, the quality of food resources drives the cannibalism of *T. lepturus*, indicating a kind of prey selectivity rather than an opportunistic feeding behaviour. The intraspecific trophic relationship is important to maintain the population of this species locally. In this area, the regular assessment of the feeding habits of this species will be important to verify possible changes in its preferred prey, which could be indicative of changes in its population density.

### 5. Acknowledgments

The author receives research grants from Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq (grant 301405/13-1) and Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro - FAPERJ (grant E-26/201.161/2014).

### 6. References

1. Baras E, Dugu R, Legendre M. Do cannibalistic fish forage optimally? An experimental study of prey size preference, bioenergetics of cannibalism and their ontogenetic variations in the African catfish *Heterobranchus longifilis*. *Aquatic Living Resources* 2014; 27:51-62.
2. Bittar VT, Castello BFL, Di Benedetto APM. Hábito alimentar do peixe-espada adulto, *Trichiurus lepturus*, na costa norte do Rio de Janeiro, sudeste do Brasil. *Biotemas* 2008; 21:83-90.
3. Bittar VT, Awabdi DR, Tonini WCT, Vidal Júnior MV, Di Benedetto APM. Feeding preference of adult females of ribbonfish *Trichiurus lepturus* L. 1758 through prey proximate-composition and caloric values. *Neotropical Ichthyology* 2012; 10:193-203.
4. Chiou WD, Chen CY, Wang CM, Chen CT. Food and feeding habits of ribbonfish *Trichiurus lepturus* in coastal waters of south-western Taiwan. *Fishery Science* 2006; 72:373-81.
5. Cruz-Torres J, Martínez-Pérez JA, Franco-Ló J, Ramírez-Villalobos AJ. Biological and ecological aspects of *Trichiurus lepturus* Linnaeus, 1758 (Perciformes: Trichiuridae) in Boca Del Rio, Veracruz, Mexico. *American-Eurasian Journal of Agriculture & Environmental Science* 2014; 14:1058-1066.
6. Di Benedetto APM, Bittar VT, Rezende CE, Camargo PB, Kehrig HA. Mercury and stable isotopes ( $^{15}\text{N}$  and  $^{13}\text{C}$ ) as tracers during the ontogeny of *Trichiurus lepturus*. *Neotropical Ichthyology* 2013; 11:211-216.
7. Di Benedetto APM, Ramos RMA. Biology of the boto-cinza dolphin (*Sotalia fluviatilis*) in south-eastern Brazil. *Journal of the Marine Biological Association of the United Kingdom* 2004; 84:1245-1250.
8. Di Benedetto APM, Siciliano S. Stomach contents of the marine tucuxi dolphin (*Sotalia guianensis*) from Rio de Janeiro, south-eastern Brazil. *Journal of the Marine Biological Association of the United Kingdom* 2007; 87:253-254.
9. Fundação Instituto de Pesca do Estado do Rio de Janeiro - FIPERJ. Diagnóstico da pesca do Estado do Rio de Janeiro, 2011. Report available in: <http://www.fiperj.rj.gov.br/index.php/publicacao/index/1>
10. Froese R, Pauly D. Fish Base, 2015. <http://www.fishbase.org>. 15 Feb, 2015.
11. Liu Y, Cheng J, Chen Y. A spatial analysis of trophic composition: a case study of hairtail (*Trichiurus japonicus*) in the East China Sea. *Hydrobiologia* 2009; 632:79-90.
12. Martins AS, Haimovici, Palacios R. Diet and feeding of the cutlassfish *Trichiurus lepturus* in the Subtropical Convergence Ecosystem of southern Brazil. *Journal of the Marine Biological Association of the United Kingdom* 2005; 85:1223-1229.
13. Prikrýl T, Novosad B. Direct evidence of cannibalism in the Oligocene cutlassfish *Anachelum glarisianum* Blainvillei, 1818 (Perciformes: Trichiuridae). *Bulletin of*

- Geosciences 2009; 84:569-572.
14. Reuben S, Vijayakumaran K, Achayya P, Prabhakar RVD. Biology and exploitation of *Trichiurus lepturus* Linnaeus from Visakhapatnam Waters. Indian Journal of Fishery 1997; 44:101-110.
  15. Smith C, Reay P. Cannibalism in teleost fish. Review of Fish Biology and Fisheries 1991; 1:41-64.
  16. Yan Y, Gang H, Chen J, Lu H, Jin X. Feeding ecology of hairtail *Trichiurus margarites* and largehead hairtail *Trichiurus lepturus* in the Beibu Gulf, the South China Sea. Chinese Journal of Oceanology and Limnology 2011; 29:174-183.